

The International Association for the Properties of Water and Steam

Moscow, Russia

June 2014

Revised Supplementary Release on Backward Equations for Specific Volume as a Function of Pressure and Temperature $v(p,T)$ for Region 3 of the IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam

© 2014 International Association for the Properties of Water and Steam
Publication in whole or in part is allowed in all countries provided that attribution is given to
the International Association for the Properties of Water and Steam

President:

Professor Tamara Petrova
Moscow Power Engineering Institute
Moscow, Russia

Executive Secretary:

Dr. R. B. Dooley
Structural Integrity Associates
Southport, Merseyside, UK
Email: bdooley@structint.com

This supplementary release replaces the corresponding supplementary release of 2005, and contains
35 pages, including this cover page.

This revised supplementary release has been authorized by the International Association for the Properties of Water and Steam (IAPWS) at its meeting in Moscow, Russia, 22-27 June, 2014, for issue by its Secretariat. The members of IAPWS are: Britain and Ireland, Canada, the Czech Republic, Germany, Japan, Russia, Scandinavia (Denmark, Finland, Norway, Sweden), and the United States, and associate members Argentina & Brazil, Australia, France, Greece, Italy, New Zealand, and Switzerland.

The backward equations $v(p,T)$ for Region 3 provided in this release are recommended as a supplement to "The IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam" (IAPWS-IF97) [1, 2]. Further details concerning the equations of this revised supplementary release can be found in the corresponding article by H.-J. Kretzschmar *et al.* [3].

This revision consists of edits to clarify descriptions of how to determine the region or subregion; the property calculations are unchanged.

Further information concerning this supplementary release, other releases, supplementary releases, guidelines, technical guidance documents, and advisory notes issued by IAPWS can be obtained from the Executive Secretary of IAPWS or from <http://www.iapws.org>.

Contents

| | |
|--|-----------|
| 1 Nomenclature | 3 |
| 2 Background | 4 |
| 3 Numerical Consistency Requirements | 5 |
| 4 Structure of the Equation Set | 5 |
| 5 Backward Equations $\nu(p,T)$ for the Subregions 3a to 3t | 6 |
| 5.1 Subregions | 6 |
| 5.2 Backward Equations $\nu(p,T)$ for the Subregions 3a to 3t | 12 |
| 5.3 Calculation of Thermodynamic Properties with the $\nu(p,T)$ Backward Equations | 13 |
| 5.4 Numerical Consistency | 14 |
| 6 Auxiliary Equations $\nu(p,T)$ for the Region very close to the Critical Point | 17 |
| 6.1 Subregions | 17 |
| 6.2 Auxiliary Equations $\nu(p,T)$ for the Subregions 3u to 3z | 19 |
| 6.3 Numerical Consistency | 20 |
| 7 Computing Time in Relation to IAPWS-IF97 | 21 |
| 8 Application of the Backward and Auxiliary Equations $\nu(p,T)$ | 22 |
| 9 References | 22 |
| Appendix | 24 |

1 Nomenclature

Thermodynamic quantities:

| | |
|----------|--------------------------------------|
| c_p | Specific isobaric heat capacity |
| f | Specific Helmholtz free energy |
| h | Specific enthalpy |
| p | Pressure |
| s | Specific entropy |
| T | Absolute temperature ^a |
| v | Specific volume |
| w | Speed of sound |
| θ | Reduced temperature $\theta = T/T^*$ |
| π | Reduced pressure, $\pi = p/p^*$ |
| ω | Reduced volume, $\omega = v/v^*$ |
| Δ | Difference in any quantity |

Subscripts:

| | |
|----------|---|
| 1...5 | Region 1...5 |
| 3a ...3z | Subregion 3a...3z |
| 3ab | Boundary between subregions 3a, 3d and 3b, 3e |
| 3cd | Boundary between subregions 3c and 3d, 3g, 3l, 3q, 3s |
| 3ef | Boundary between subregions 3e, 3h, 3n and 3f, 3i, 3o |
| 3gh | Boundary between subregions 3g, 3l and 3h, 3m |
| 3ij | Boundary between subregions 3i, 3p and 3j |
| 3jk | Boundary between subregions 3j, 3r and 3k |
| 3mn | Boundary between subregions 3m and 3n |
| 3op | Boundary between subregions 3o and 3p |
| 3qu | Boundary between of subregion 3q and 3u |
| 3rx | Boundary between of subregion 3r and 3x |
| 3uv | Boundary between subregions 3u and 3v |
| 3wx | Boundary between subregions 3w and 3x |
| B23 | Boundary between regions 2 and 3 |
| c | Critical point |
| it | Iterated quantity |
| max | Maximum value of a quantity |
| RMS | Root-mean-square value of a quantity |
| sat | Saturation state |
| tol | Tolerated value of a quantity |

Root-mean-square value:

$$\Delta x_{\text{RMS}} = \sqrt{\frac{1}{N} \sum_{n=1}^N (\Delta x_n)^2}$$

where Δx_n can be either absolute or percentage difference between the corresponding quantities x ; N is the number of Δx_n values (10 million points uniformly distributed over the range of validity in the p - T plane).

Superscripts:

| | |
|----|------------------------------------|
| 97 | Quantity or equation of IAPWS-IF97 |
| 01 | Equation of IAPWS-IF97-S01 |
| 03 | Equation of IAPWS-IF97-S03rev |
| 04 | Equation of IAPWS-IF97-S04 |
| * | Reducing quantity |
| ' | Saturated liquid state |
| " | Saturated vapor state |

^a Note: T denotes absolute temperature on the International Temperature Scale of 1990 (ITS-90).

2 Background

The IAPWS Industrial Formulation 1997 for the thermodynamic properties of water and steam (IAPWS-IF97) [1, 2] contains basic equations, saturation equations and equations for the frequently used backward functions $T(p, h)$ and $T(p, s)$ valid in the liquid region 1 and the vapor region 2; see Figure 1. IAPWS-IF97 was supplemented by "Supplementary Release on Backward Equations for Pressure as a Function of Enthalpy and Entropy $p(h, s)$ to the IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam" [4, 5], which will be referred to as IAPWS-IF97-S01. These equations are valid in region 1 and region 2. An additional "Supplementary Release on Backward Equations for the Functions $T(p, h)$, $v(p, h)$ and $T(p, s)$, $v(p, s)$ for Region 3 of the IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam" [6, 7], which will be referred to as IAPWS-IF97-S03rev, was adopted by IAPWS in 2003 and revised in 2004. In 2004, IAPWS-IF97 was supplemented by "Supplementary Release on Backward Equations $p(h, s)$ for Region 3, Equations as a Function of h and s for the Region Boundaries, and an Equation $T_{\text{sat}}(h, s)$ for Region 4 of the IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam" (referred to here as IAPWS-IF97-S04) [8, 9].




Figure 1. Regions and equations of IAPWS-IF97, IAPWS-IF97-S01, IAPWS-IF97-S03rev, IAPWS-IF97-S04, and the equations $v_3(p, T)$ of this release

IAPWS-IF97 region 3 is covered by a basic equation for the Helmholtz free energy $f(v, T)$. All thermodynamic properties can be derived from the basic equation as a function of specific volume v and temperature T . However, in modeling some steam power cycles, thermodynamic properties as functions of the variables (p, T) are required in region 3. It is cumbersome to perform these calculations with IAPWS-IF97, because they require iterations of v from p and T using the function $p(v, T)$ derived from the IAPWS-IF97 basic equation $f(v, T)$.

In order to avoid such iterations, this release provides equations $v_3(p, T)$; see Figure 1. With specific volume v calculated from the equations $v_3(p, T)$, the other properties in region 3 can be calculated using the IAPWS-IF97 basic equation $f(v, T)$.

For process calculations, the numerical consistency requirements for the equations $v(p, T)$ are very strict. Because the specific volume in the p - T plane has a complicated structure, including an infinite slope at the critical point, region 3 was divided into 26 subregions. The first 20 subregions and their associated backward equations, described in Section 5, cover almost all of region 3 and fully meet the consistency requirements. For a small area very near the critical point, it was not possible to meet the consistency requirements fully. This near-critical region is covered with reasonable consistency by six subregions with auxiliary equations, described in Section 6.

3 Numerical Consistency Requirements

The permissible value for the numerical consistency of the equations for specific volume with the IAPWS-IF97 fundamental equation was determined based on the required accuracy of the iteration otherwise used. The iteration accuracy depends on thermodynamic process calculations. To obtain specific enthalpy or entropy from pressure and temperature in region 3 with a maximum deviation of 0.001 % from IAPWS-IF97, and isobaric heat capacity or speed of sound with a maximum deviation of 0.01 %, a relative accuracy of $|\Delta v/v| = 0.001\%$ is sufficient. Therefore, the permissible relative tolerance for the equations $v(p, T)$ was set to $|\Delta v/v|_{\text{tol}} = 0.001\%$.

4 Structure of the Equation Set

The range of validity of the equations $v_3(p, T)$ is region 3 defined by:

$$623.15 \text{ K} < T \leq 863.15 \text{ K} \text{ and } p_{\text{B23}}^{97}(T) < p \leq 100 \text{ MPa.}$$

The function $p_{\text{B23}}^{97}(T)$ represents the B23-equation of IAPWS-IF97.

It proved to be infeasible to achieve the numerical consistency requirement of 0.001 % for $v_3(p, T)$ using simple functional forms in the region

$$T_{3\text{qu}}(p) < T \leq T_{3\text{rx}}(p) \text{ for } p_{\text{sat}}^{97}(643.15 \text{ K}) < p \leq 22.5 \text{ MPa}; \text{ see Figure 2.}$$

This limitation is due to the infinite slope of the specific volume at the critical point. In order to cover region 3 completely, Section 6 contains auxiliary equations for this small region very close to the critical point.

Figure 2 shows the range of validity of the backward and auxiliary equations.




Figure 2. Range of validity of the backward and auxiliary equations. The area marked in gray is not true to scale but enlarged to make the small area better visible.

5 Backward Equations $v(p, T)$ for the Subregions 3a to 3t

5.1 Subregions

Preliminary investigations showed that it was not possible to meet the numerical consistency requirement with only a few $v(p, T)$ equations. Therefore, the main part of region 3 was divided into 20 subregions 3a to 3t; see Figures 3 and 4.




Figure 3. Division of region 3 into subregions for the backward equations $v_3(p, T)$




Figure 4. Enlargement from Figure 3 for the subregions 3c to 3r for the backward equation $v(p, T)$

The subregion boundary equations, except for $T_{3ab}(p)$, $T_{3ef}(p)$, and $T_{3op}(p)$, have the following dimensionless form:

$$\frac{T(p)}{T^*} = \theta(\pi) = \sum_{i=1}^N n_i \pi^{I_i} , \quad (1)$$

where $\theta = T/T^*$, $\pi = p/p^*$, with $T^* = 1 \text{ K}$, $p^* = 1 \text{ MPa}$.

The equations $T_{3ab}(p)$ and $T_{3op}(p)$ have the form:

$$\frac{T(p)}{T^*} = \theta(\pi) = \sum_{i=1}^N n_i (\ln \pi)^{I_i} , \quad (2)$$

and $T_{3ef}(p)$ has the form:

$$\frac{T_{3ef}(p)}{T^*} = \theta_{3ef}(\pi) = \left. \frac{\partial \theta_{\text{sat}}}{\partial \pi} \right|_c (\pi - 22.064) + 647.096 , \quad (3)$$

where $\left. \frac{\partial \theta_{\text{sat}}}{\partial \pi} \right|_c = 3.727 \ 888 \ 004$.

The coefficients n_i and the exponents I_i of the boundary equations are listed in Table 1.

Table 1. Numerical values of the coefficients of the equations for subregion boundaries (except $T_{3ef}(p)$)

| Equation | i | I_i | n_i | i | I_i | n_i |
|--------------|-----|-------|--------------------------------------|-----|-------|---|
| $T_{3ab}(p)$ | 1 | 0 | 0.154 793 642 129 415 $\times 10^4$ | 4 | -1 | -0.191 887 498 864 292 $\times 10^4$ |
| | 2 | 1 | -0.187 661 219 490 113 $\times 10^3$ | 5 | -2 | 0.918 419 702 359 447 $\times 10^3$ |
| | 3 | 2 | 0.213 144 632 222 113 $\times 10^2$ | | | |
| $T_{3cd}(p)$ | 1 | 0 | 0.585 276 966 696 349 $\times 10^3$ | 3 | 2 | -0.127 283 549 295 878 $\times 10^{-1}$ |
| | 2 | 1 | 0.278 233 532 206 915 $\times 10^1$ | 4 | 3 | 0.159 090 746 562 729 $\times 10^{-3}$ |
| $T_{3gh}(p)$ | 1 | 0 | -0.249 284 240 900 418 $\times 10^5$ | 4 | 3 | 0.751 608 051 114 157 $\times 10^1$ |
| | 2 | 1 | 0.428 143 584 791 546 $\times 10^4$ | 5 | 4 | -0.787 105 249 910 383 $\times 10^{-1}$ |
| | 3 | 2 | -0.269 029 173 140 130 $\times 10^3$ | | | |
| $T_{3ij}(p)$ | 1 | 0 | 0.584 814 781 649 163 $\times 10^3$ | 4 | 3 | -0.587 071 076 864 459 $\times 10^{-2}$ |
| | 2 | 1 | -0.616 179 320 924 617 | 5 | 4 | 0.515 308 185 433 082 $\times 10^{-4}$ |
| | 3 | 2 | 0.260 763 050 899 562 | | | |
| $T_{3jk}(p)$ | 1 | 0 | 0.617 229 772 068 439 $\times 10^3$ | 4 | 3 | -0.157 391 839 848 015 $\times 10^{-1}$ |
| | 2 | 1 | -0.770 600 270 141 675 $\times 10^1$ | 5 | 4 | 0.137 897 492 684 194 $\times 10^{-3}$ |
| | 3 | 2 | 0.697 072 596 851 896 | | | |
| $T_{3mn}(p)$ | 1 | 0 | 0.535 339 483 742 384 $\times 10^3$ | 3 | 2 | -0.158 365 725 441 648 |
| | 2 | 1 | 0.761 978 122 720 128 $\times 10^1$ | 4 | 3 | 0.192 871 054 508 108 $\times 10^{-2}$ |
| $T_{3op}(p)$ | 1 | 0 | 0.969 461 372 400 213 $\times 10^3$ | 4 | -1 | 0.773 845 935 768 222 $\times 10^3$ |
| | 2 | 1 | -0.332 500 170 441 278 $\times 10^3$ | 5 | -2 | -0.152 313 732 937 084 $\times 10^4$ |
| | 3 | 2 | 0.642 859 598 466 067 $\times 10^2$ | | | |
| $T_{3qu}(p)$ | 1 | 0 | 0.565 603 648 239 126 $\times 10^3$ | 3 | 2 | -0.102 020 639 611 016 |
| | 2 | 1 | 0.529 062 258 221 222 $\times 10^1$ | 4 | 3 | 0.122 240 301 070 145 $\times 10^{-2}$ |
| $T_{3rx}(p)$ | 1 | 0 | 0.584 561 202 520 006 $\times 10^3$ | 3 | 2 | 0.243 293 362 700 452 |
| | 2 | 1 | -0.102 961 025 163 669 $\times 10^1$ | 4 | 3 | -0.294 905 044 740 799 $\times 10^{-2}$ |

The following description of the use of the subregion boundary equations is summarized in Table 2 and Figures 3 and 4.

Table 2. Pressure ranges and corresponding subregion boundary equations for determining the correct subregion, 3a to 3t, for the backward equations $v(p, T)$

| Pressure Range | Sub-region | For | Sub-region | For |
|---|------------|--|------------|---|
| $40 \text{ MPa} < p \leq 100 \text{ MPa}$ | 3a | $T \leq T_{3ab}(p)$ | 3b | $T > T_{3ab}(p)$ |
| $25 \text{ MPa} < p \leq 40 \text{ MPa}$ | 3c | $T \leq T_{3cd}(p)$ | 3e | $T_{3ab}(p) < T \leq T_{3ef}(p)$ |
| | 3d | $T_{3cd}(p) < T \leq T_{3ab}(p)$ | 3f | $T > T_{3ef}(p)$ |
| $23.5 \text{ MPa} < p \leq 25 \text{ MPa}$ | 3c | $T \leq T_{3cd}(p)$ | 3i | $T_{3ef}(p) < T \leq T_{3ij}(p)$ |
| | 3g | $T_{3cd}(p) < T \leq T_{3gh}(p)$ | 3j | $T_{3ij}(p) < T \leq T_{3jk}(p)$ |
| | 3h | $T_{3gh}(p) < T \leq T_{3ef}(p)$ | 3k | $T > T_{3jk}(p)$ |
| $23 \text{ MPa} < p \leq 23.5 \text{ MPa}$ | 3c | $T \leq T_{3cd}(p)$ | 3i | $T_{3ef}(p) < T \leq T_{3ij}(p)$ |
| | 3l | $T_{3cd}(p) < T \leq T_{3gh}(p)$ | 3j | $T_{3ij}(p) < T \leq T_{3jk}(p)$ |
| | 3h | $T_{3gh}(p) < T \leq T_{3ef}(p)$ | 3k | $T > T_{3jk}(p)$ |
| $22.5 \text{ MPa} < p \leq 23 \text{ MPa}$ | 3c | $T \leq T_{3cd}(p)$ | 3o | $T_{3ef}(p) < T \leq T_{3op}(p)$ |
| | 3l | $T_{3cd}(p) < T \leq T_{3gh}(p)$ | 3p | $T_{3op}(p) < T \leq T_{3ij}(p)$ |
| | 3m | $T_{3gh}(p) < T \leq T_{3mn}(p)$ | 3j | $T_{3ij}(p) < T \leq T_{3jk}(p)$ |
| | 3n | $T_{3mn}(p) < T \leq T_{3ef}(p)$ | 3k | $T > T_{3jk}(p)$ |
| $p_{\text{sat}}^{97}(643.15 \text{ K}) < p \leq 22.5 \text{ MPa}$ | 3c | $T \leq T_{3cd}(p)$ | 3r | $T_{3rx}(p) < T \leq T_{3ik}(p)$ |
| | 3q | $T_{3cd}(p) < T \leq T_{3qu}(p)$ | 3k | $T > T_{3jk}(p)$ |
| $20.5 \text{ MPa} < p \leq p_{\text{sat}}^{97}(643.15 \text{ K})$ | 3c | $T \leq T_{3cd}(p)$ | 3r | $T_{\text{sat}}^{97}(p) \leq T \leq T_{3jk}(p)$ |
| | 3s | $T_{3cd}(p) < T \leq T_{\text{sat}}^{97}(p)$ | 3k | $T > T_{3jk}(p)$ |
| $p_{3cd}^b < p \leq 20.5 \text{ MPa}$ | 3c | $T \leq T_{3cd}(p)$ | 3t | $T \geq T_{\text{sat}}^{97}(p)$ |
| | 3s | $T_{3cd}(p) < T \leq T_{\text{sat}}^{97}(p)$ | | |
| $p_{\text{sat}}^{97}(623.15 \text{ K}) < p \leq p_{3cd}^b$ | 3c | $T \leq T_{\text{sat}}^{97}(p)$ | 3t | $T \geq T_{\text{sat}}^{97}(p)$ |

$${}^b p_{3cd} = 1.900\ 881\ 189\ 173\ 929 \times 10^1 \text{ MPa}$$

The equation $T_{3ab}(p)$ approximates the critical isentrope from 25 MPa to 100 MPa and represents the boundary equation between subregion 3a and subregion 3d.

The equation $T_{3cd}(p)$ ranges from $p_{3cd} = 1.900\ 881\ 189\ 173\ 929 \times 10^1 \text{ MPa}$ to 40 MPa. The pressure of $p_{3cd} = 1.900\ 881\ 189\ 173\ 929 \times 10^1 \text{ MPa}$ is given as $T_{\text{sat}}^{97}(p) - T_{3cd}(p) = 0$. The equation $T_{3cd}(p)$ represents the boundary equation between subregions 3d, 3g, 3l, 3q or 3s, and subregion 3c.

The equation $T_{3gh}(p)$ ranges from 22.5 MPa to 25 MPa and represents the boundary equation between subregions 3h or 3m and subregions 3g or 3l.

The equation $T_{3ij}(p)$ approximates the isochore $v = 0.0041 \text{ m}^3 \text{ kg}^{-1}$ from 22.5 MPa to 25 MPa and represents the boundary equation between subregion 3j and subregions 3i or 3p.

The equation $T_{3jk}(p)$ approximates the isochore $v = v''(20.5 \text{ MPa})$ from 20.5 MPa to 25 MPa and represents the boundary equation between subregion 3k and subregions 3j or 3r.

The equation $T_{3mn}(p)$ approximates the isochore $v = 0.0028 \text{ m}^3 \text{ kg}^{-1}$ from 22.5 MPa to 23 MPa and represents the boundary equation between subregion 3n and subregion 3m.

The equation $T_{3op}(p)$ approximates the isochore $v = 0.0034 \text{ m}^3 \text{ kg}^{-1}$ from 22.5 MPa to 23 MPa and represents the boundary equation between subregion 3p and subregion 3o.

The equation $T_{3qu}(p)$ approximates the isochore $v = v'(643.15 \text{ K})$ from $p = p_{\text{sat}}^{97}(643.15 \text{ K})$, where $p_{\text{sat}}^{97}(643.15 \text{ K}) = 2.104\,336\,732 \times 10^1 \text{ MPa}$ to 22.5 MPa and represents the boundary equation between subregion 3q and subregion 3r (see Fig. 5).

The equation $T_{3rx}(p)$ approximates the isochore $v = v''(643.15 \text{ K})$ from $p = p_{\text{sat}}^{97}(643.15 \text{ K})$, where $p_{\text{sat}}^{97}(643.15 \text{ K}) = 2.104\,336\,732 \times 10^1 \text{ MPa}$, to 22.5 MPa and represents the boundary equation between subregion 3r and subregion 3x (see Fig.5).

The subregion boundary equation $T_{3ef}(p)$ is a straight line from 22.064 MPa to 40 MPa having the slope of the saturation-temperature curve of IAPWS-IF97 at the critical point. It divides subregions 3f, 3i or 3o from subregions 3e, 3h or 3n.

Computer-program verification

To assist the user in computer-program verification of the equations for the subregion boundaries, Table 3 contains test values for calculated temperatures.

Table 3. Selected temperature values calculated from the subregion boundary equations^c

| Equation | p MPa | T K | Equation | p MPa | T K |
|--------------|------------|-------------------------------|--------------|------------|-------------------------------|
| $T_{3ab}(p)$ | 40 | $6.930\,341\,408 \times 10^2$ | $T_{3jk}(p)$ | 23 | $6.558\,338\,344 \times 10^2$ |
| $T_{3cd}(p)$ | 25 | $6.493\,659\,208 \times 10^2$ | $T_{3mn}(p)$ | 22.8 | $6.496\,054\,133 \times 10^2$ |
| $T_{3ef}(p)$ | 40 | $7.139\,593\,992 \times 10^2$ | $T_{3op}(p)$ | 22.8 | $6.500\,106\,943 \times 10^2$ |
| $T_{3gh}(p)$ | 23 | $6.498\,873\,759 \times 10^2$ | $T_{3qu}(p)$ | 22 | $6.456\,355\,027 \times 10^2$ |
| $T_{3ij}(p)$ | 23 | $6.515\,778\,091 \times 10^2$ | $T_{3rx}(p)$ | 22 | $6.482\,622\,754 \times 10^2$ |

^c It is recommended that programmed functions be verified using 8 byte real values for all variables.

5.2 Backward Equations $v(p,T)$ for the Subregions 3a to 3t

The backward equations $v(p,T)$ for the subregions 3a to 3t, except for 3n, have the following dimensionless form:

$$\frac{v(p,T)}{v^*} = \omega(\pi, \theta) = \left[\sum_{i=1}^N n_i [(\pi - a)^c]^{I_i} [(\theta - b)^d]^{J_i} \right]^e. \quad (4)$$

The equation for subregion 3n has the form:

$$\frac{v_{3n}(p,T)}{v^*} = \omega_{3n}(\pi, \theta) = \exp \left\{ \sum_{i=1}^N n_i (\pi - a)^{I_i} (\theta - b)^{J_i} \right\}, \quad (5)$$

with $\omega = v/v^*$, $\pi = p/p^*$, and $\theta = T/T^*$. The reducing quantities v^* , p^* , and T^* , the number of coefficients N , the non-linear parameters a and b , and the exponents c , d , and e are listed in Table 4 for the equations of the subregions 3a to 3t. The coefficients n_i and exponents I_i and J_i of these equations are listed in Tables A1.1 to A1.20 of the Appendix.

Table 4. Reducing quantities v^* , p^* , and T^* , number of coefficients N , non-linear parameters a and b , and exponents c , d , and e for the $v(p,T)$ equations of the subregions 3a to 3t

| Subregion | v^* $\text{m}^3 \text{ kg}^{-1}$ | p^* MPa | T^* K | N | a | b | c | d | e |
|-----------|---------------------------------------|--------------|------------|-----|-------|-------|-----|------|-----|
| 3a | 0.0024 | 100 | 760 | 30 | 0.085 | 0.817 | 1 | 1 | 1 |
| 3b | 0.0041 | 100 | 860 | 32 | 0.280 | 0.779 | 1 | 1 | 1 |
| 3c | 0.0022 | 40 | 690 | 35 | 0.259 | 0.903 | 1 | 1 | 1 |
| 3d | 0.0029 | 40 | 690 | 38 | 0.559 | 0.939 | 1 | 1 | 4 |
| 3e | 0.0032 | 40 | 710 | 29 | 0.587 | 0.918 | 1 | 1 | 1 |
| 3f | 0.0064 | 40 | 730 | 42 | 0.587 | 0.891 | 0.5 | 1 | 4 |
| 3g | 0.0027 | 25 | 660 | 38 | 0.872 | 0.971 | 1 | 1 | 4 |
| 3h | 0.0032 | 25 | 660 | 29 | 0.898 | 0.983 | 1 | 1 | 4 |
| 3i | 0.0041 | 25 | 660 | 42 | 0.910 | 0.984 | 0.5 | 1 | 4 |
| 3j | 0.0054 | 25 | 670 | 29 | 0.875 | 0.964 | 0.5 | 1 | 4 |
| 3k | 0.0077 | 25 | 680 | 34 | 0.802 | 0.935 | 1 | 1 | 1 |
| 3l | 0.0026 | 24 | 650 | 43 | 0.908 | 0.989 | 1 | 1 | 4 |
| 3m | 0.0028 | 23 | 650 | 40 | 1.00 | 0.997 | 1 | 0.25 | 1 |
| 3n | 0.0031 | 23 | 650 | 39 | 0.976 | 0.997 | - | - | - |
| 3o | 0.0034 | 23 | 650 | 24 | 0.974 | 0.996 | 0.5 | 1 | 1 |
| 3p | 0.0041 | 23 | 650 | 27 | 0.972 | 0.997 | 0.5 | 1 | 1 |
| 3q | 0.0022 | 23 | 650 | 24 | 0.848 | 0.983 | 1 | 1 | 4 |
| 3r | 0.0054 | 23 | 650 | 27 | 0.874 | 0.982 | 1 | 1 | 1 |
| 3s | 0.0022 | 21 | 640 | 29 | 0.886 | 0.990 | 1 | 1 | 4 |
| 3t | 0.0088 | 20 | 650 | 33 | 0.803 | 1.02 | 1 | 1 | 1 |

Computer-program verification

To assist the user in computer-program verification of the equations for the subregions 3a to 3t, Table 5 contains test values for calculated specific volumes.

Table 5. Selected specific volume values calculated from the equations for the subregions 3a to 3t^d

| Equation | p MPa | T K | v $\text{m}^3 \text{kg}^{-1}$ | Equation | p MPa | T K | v $\text{m}^3 \text{kg}^{-1}$ |
|---------------|------------|----------|------------------------------------|---------------|------------|----------|------------------------------------|
| $v_{3a}(p,T)$ | 50 | 630 | $1.470\ 853\ 100 \times 10^{-3}$ | $v_{3k}(p,T)$ | 23 | 660 | $6.109\ 525\ 997 \times 10^{-3}$ |
| | 80 | 670 | $1.503\ 831\ 359 \times 10^{-3}$ | | 24 | 670 | $6.427\ 325\ 645 \times 10^{-3}$ |
| $v_{3b}(p,T)$ | 50 | 710 | $2.204\ 728\ 587 \times 10^{-3}$ | $v_{3l}(p,T)$ | 22.6 | 646 | $2.117\ 860\ 851 \times 10^{-3}$ |
| | 80 | 750 | $1.973\ 692\ 940 \times 10^{-3}$ | | 23 | 646 | $2.062\ 374\ 674 \times 10^{-3}$ |
| $v_{3c}(p,T)$ | 20 | 630 | $1.761\ 696\ 406 \times 10^{-3}$ | $v_{3m}(p,T)$ | 22.6 | 648.6 | $2.533\ 063\ 780 \times 10^{-3}$ |
| | 30 | 650 | $1.819\ 560\ 617 \times 10^{-3}$ | | 22.8 | 649.3 | $2.572\ 971\ 781 \times 10^{-3}$ |
| $v_{3d}(p,T)$ | 26 | 656 | $2.245\ 587\ 720 \times 10^{-3}$ | $v_{3n}(p,T)$ | 22.6 | 649.0 | $2.923\ 432\ 711 \times 10^{-3}$ |
| | 30 | 670 | $2.506\ 897\ 702 \times 10^{-3}$ | | 22.8 | 649.7 | $2.913\ 311\ 494 \times 10^{-3}$ |
| $v_{3e}(p,T)$ | 26 | 661 | $2.970\ 225\ 962 \times 10^{-3}$ | $v_{3o}(p,T)$ | 22.6 | 649.1 | $3.131\ 208\ 996 \times 10^{-3}$ |
| | 30 | 675 | $3.004\ 627\ 086 \times 10^{-3}$ | | 22.8 | 649.9 | $3.221\ 160\ 278 \times 10^{-3}$ |
| $v_{3f}(p,T)$ | 26 | 671 | $5.019\ 029\ 401 \times 10^{-3}$ | $v_{3p}(p,T)$ | 22.6 | 649.4 | $3.715\ 596\ 186 \times 10^{-3}$ |
| | 30 | 690 | $4.656\ 470\ 142 \times 10^{-3}$ | | 22.8 | 650.2 | $3.664\ 754\ 790 \times 10^{-3}$ |
| $v_{3g}(p,T)$ | 23.6 | 649 | $2.163\ 198\ 378 \times 10^{-3}$ | $v_{3q}(p,T)$ | 21.1 | 640 | $1.970\ 999\ 272 \times 10^{-3}$ |
| | 24 | 650 | $2.166\ 044\ 161 \times 10^{-3}$ | | 21.8 | 643 | $2.043\ 919\ 161 \times 10^{-3}$ |
| $v_{3h}(p,T)$ | 23.6 | 652 | $2.651\ 081\ 407 \times 10^{-3}$ | $v_{3r}(p,T)$ | 21.1 | 644 | $5.251\ 009\ 921 \times 10^{-3}$ |
| | 24 | 654 | $2.967\ 802\ 335 \times 10^{-3}$ | | 21.8 | 648 | $5.256\ 844\ 741 \times 10^{-3}$ |
| $v_{3i}(p,T)$ | 23.6 | 653 | $3.273\ 916\ 816 \times 10^{-3}$ | $v_{3s}(p,T)$ | 19.1 | 635 | $1.932\ 829\ 079 \times 10^{-3}$ |
| | 24 | 655 | $3.550\ 329\ 864 \times 10^{-3}$ | | 20 | 638 | $1.985\ 387\ 227 \times 10^{-3}$ |
| $v_{3j}(p,T)$ | 23.5 | 655 | $4.545\ 001\ 142 \times 10^{-3}$ | $v_{3t}(p,T)$ | 17 | 626 | $8.483\ 262\ 001 \times 10^{-3}$ |
| | 24 | 660 | $5.100\ 267\ 704 \times 10^{-3}$ | | 20 | 640 | $6.227\ 528\ 101 \times 10^{-3}$ |

^d It is recommended that programmed functions be verified using 8 byte real values for all variables.

5.3 Calculation of Thermodynamic Properties with the $v(p,T)$ Backward Equations

The $v(p,T)$ backward equations described in Section 5.2 together with IAPWS-IF97 basic equation $f(v,T)$ make it possible to determine all thermodynamic properties, *e.g.*, enthalpy, entropy, isobaric heat capacity, speed of sound, from pressure p and temperature T in region 3 without iteration.

The following steps should be made:

- Identify the subregion (3a to 3t) for given pressure p and temperature T following the instructions of Section 5.1 in conjunction with Table 2 and Figures 3 and 4. Then, calculate the specific volume v for the subregion using the corresponding backward equation $v(p,T)$.
- Calculate the desired thermodynamic property from the previously calculated specific volume v and the given temperature T using the derivatives of the IAPWS-IF97 basic equation $f(v,T)$, where $v = v(p,T)$; see Table 31 in [1].

5.4 Numerical Consistency

5.4.1 Numerical Consistency with the Basic Equation of IAPWS-IF97

The maximum relative deviations and root-mean-square relative deviations of specific volume, calculated from the backward equations $v(p,T)$ for subregions 3a to 3t, from the IAPWS-IF97 basic equation $f(v,T)$ in comparison with the permissible tolerances are listed in Table 6. The calculation of the root-mean-square values is described in Section 1.

Table 6 also contains the maximum relative deviations and root-mean-square relative deviations of specific enthalpy, specific entropy, specific isobaric heat capacity, and speed of sound, calculated as described in Section 5.3.

Table 6. Maximum relative deviations and root-mean-square relative deviations of the specific volume, calculated from the backward equations for subregions 3a to 3t, and maximum relative deviations of specific enthalpy, specific entropy, specific isobaric heat capacity and speed of sound, calculated as described in Section 5.3, from the IAPWS-IF97 basic equation $f(v,T)$

| Subregion | $ \Delta v/v $ | | $ \Delta h/h $ | | $ \Delta s/s $ | | $ \Delta c_p/c_p $ | | $ \Delta w/w $ | |
|-----------------------|----------------|---------|----------------|---------|----------------|---------|--------------------|--------|----------------|--------|
| | % | | % | | % | | % | | % | |
| | max | RMS | max | RMS | max | RMS | max | RMS | max | RMS |
| 3a | 0.00061 | 0.00031 | 0.00018 | 0.00008 | 0.00026 | 0.00011 | 0.0016 | 0.0006 | 0.0015 | 0.0006 |
| 3b | 0.00064 | 0.00035 | 0.00017 | 0.00008 | 0.00016 | 0.00008 | 0.0012 | 0.0003 | 0.0008 | 0.0003 |
| 3c | 0.00080 | 0.00038 | 0.00026 | 0.00012 | 0.00025 | 0.00011 | 0.0059 | 0.0016 | 0.0023 | 0.0010 |
| 3d | 0.00059 | 0.00025 | 0.00018 | 0.00008 | 0.00014 | 0.00006 | 0.0035 | 0.0010 | 0.0012 | 0.0004 |
| 3e | 0.00072 | 0.00033 | 0.00018 | 0.00009 | 0.00014 | 0.00007 | 0.0017 | 0.0005 | 0.0006 | 0.0002 |
| 3f | 0.00068 | 0.00020 | 0.00018 | 0.00005 | 0.00013 | 0.00004 | 0.0015 | 0.0003 | 0.0002 | 0.0001 |
| 3g | 0.00047 | 0.00016 | 0.00014 | 0.00005 | 0.00011 | 0.00004 | 0.0032 | 0.0011 | 0.0010 | 0.0003 |
| 3h | 0.00085 | 0.00044 | 0.00022 | 0.00012 | 0.00017 | 0.00009 | 0.0066 | 0.0018 | 0.0006 | 0.0002 |
| 3i | 0.00067 | 0.00028 | 0.00018 | 0.00008 | 0.00013 | 0.00006 | 0.0019 | 0.0006 | 0.0002 | 0.0001 |
| 3j | 0.00034 | 0.00019 | 0.00009 | 0.00005 | 0.00007 | 0.00004 | 0.0020 | 0.0006 | 0.0002 | 0.0001 |
| 3k | 0.00034 | 0.00012 | 0.00008 | 0.00003 | 0.00007 | 0.00002 | 0.0018 | 0.0003 | 0.0002 | 0.0001 |
| 3l | 0.00033 | 0.00019 | 0.00010 | 0.00006 | 0.00008 | 0.00005 | 0.0035 | 0.0015 | 0.0008 | 0.0004 |
| 3m | 0.00057 | 0.00031 | 0.00015 | 0.00009 | 0.00011 | 0.00006 | 0.0062 | 0.0030 | 0.0006 | 0.0002 |
| 3n | 0.00064 | 0.00029 | 0.00017 | 0.00008 | 0.00012 | 0.00006 | 0.0050 | 0.0013 | 0.0002 | 0.0001 |
| 3o | 0.00031 | 0.00015 | 0.00008 | 0.00004 | 0.00006 | 0.00003 | 0.0007 | 0.0002 | 0.0001 | 0.0001 |
| 3p | 0.00044 | 0.00022 | 0.00012 | 0.00006 | 0.00009 | 0.00005 | 0.0026 | 0.0010 | 0.0002 | 0.0001 |
| 3q | 0.00036 | 0.00018 | 0.00012 | 0.00006 | 0.00009 | 0.00005 | 0.0040 | 0.0016 | 0.0010 | 0.0005 |
| 3r | 0.00037 | 0.00007 | 0.00010 | 0.00002 | 0.00008 | 0.00002 | 0.0030 | 0.0004 | 0.0002 | 0.0001 |
| 3s | 0.00030 | 0.00016 | 0.00010 | 0.00005 | 0.00007 | 0.00004 | 0.0033 | 0.0015 | 0.0009 | 0.0005 |
| 3t | 0.00095 | 0.00045 | 0.00022 | 0.00010 | 0.00018 | 0.00008 | 0.0046 | 0.0015 | 0.0004 | 0.0002 |
| permissible tolerance | 0.001 | | 0.001 | | 0.001 | | 0.01 | | 0.01 | |

Table 6 shows that the deviations of the specific volume, specific enthalpy, and specific entropy from the IAPWS-IF97 basic equation are less than 0.001 % and the deviations of specific isobaric heat capacity and speed of sound are less than 0.01 %. Therefore, the values

of specific volume, specific enthalpy and specific entropy of IAPWS-IF97 are represented with 5 significant figures, and the values of specific isobaric heat capacity and speed of sound with 4 significant figures by using the backward equations $v(p, T)$.

5.4.2 Consistency at Boundaries Between Subregions

The maximum relative differences of specific volume between the $v(p, T)$ backward equations of adjacent subregions along the subregion boundary pressures are listed in the third column of Table 7. Table 8 contains these maximum relative differences along the subregion boundary equations.

Table 7. Maximum relative deviations of specific volume between the backward equations $v(p, T)$ of adjacent subregions and maximum relative deviations of specific enthalpy, specific entropy, specific isobaric heat capacity, and speed of sound, calculated as described in Section 5.3, along the subregion boundary pressures

| Subregion Boundary | Between Subregions | $ \Delta v/v _{\max}$ % | $ \Delta h/h _{\max}$ % | $ \Delta s/s _{\max}$ % | $ \Delta c_p/c_p _{\max}$ % | $ \Delta w/w _{\max}$ % |
|---|--------------------|----------------------------|----------------------------|----------------------------|--------------------------------|----------------------------|
| $p = 40$ MPa | 3a, 3c | 0.00074 | 0.00021 | 0.00028 | 0.0018 | 0.0019 |
| | 3a, 3d | 0.00060 | 0.00017 | 0.00013 | 0.0013 | 0.0006 |
| | 3b, 3e | 0.00062 | 0.00015 | 0.00012 | 0.0009 | 0.0004 |
| | 3b, 3f | 0.00078 | 0.00018 | 0.00014 | 0.0004 | 0.0002 |
| $p = 25$ MPa | 3d, 3g | 0.00056 | 0.00015 | 0.00011 | 0.0031 | 0.0010 |
| | 3d, 3h | 0.00056 | 0.00015 | 0.00011 | 0.0021 | 0.0003 |
| | 3e, 3h | 0.00063 | 0.00017 | 0.00013 | 0.0014 | 0.0002 |
| | 3f, 3i | 0.00055 | 0.00014 | 0.00011 | 0.0011 | 0.0002 |
| | 3f, 3j | 0.00060 | 0.00015 | 0.00011 | 0.0015 | 0.0002 |
| | 3f, 3k | 0.00064 | 0.00013 | 0.00011 | 0.0011 | 0.0002 |
| $p = 23.5$ MPa | 3g, 3l | 0.00049 | 0.00015 | 0.00012 | 0.0033 | 0.0011 |
| $p = 23$ MPa | 3h, 3m | 0.00084 | 0.00023 | 0.00017 | 0.0074 | 0.0007 |
| | 3h, 3n | 0.00085 | 0.00022 | 0.00016 | 0.0047 | 0.0003 |
| | 3i, 3o | 0.00047 | 0.00012 | 0.00009 | 0.0006 | 0.0002 |
| | 3i, 3p | 0.00059 | 0.00015 | 0.00012 | 0.0020 | 0.0002 |
| $p = 22.5$ MPa | 3l, 3q | 0.00033 | 0.00010 | 0.00008 | 0.0025 | 0.0008 |
| | 3j, 3r | 0.00035 | 0.00009 | 0.00007 | 0.0015 | 0.0002 |
| $p = p_{\text{sat}}^{97}(643.15 \text{ K})$ | 3q, 3s | 0.00033 | 0.00010 | 0.00008 | 0.0036 | 0.0008 |
| $p = 20.5$ MPa | 3k, 3t | 0.00042 | 0.00009 | 0.00008 | 0.0019 | 0.0002 |
| permissible tolerance | | 0.001 | 0.001 | 0.001 | 0.01 | 0.01 |

Table 8. Maximum relative deviations of specific volume between the backward equations $v(p, T)$ of the adjacent subregions and maximum relative deviations of specific enthalpy, specific entropy, specific isobaric heat capacity, and speed of sound, calculated as described in Section 5.3, along the subregion boundary equations

| Subregion Boundary Equation | Between Subregions | $ \Delta v/v _{\max}$ % | $ \Delta h/h _{\max}$ % | $ \Delta s/s _{\max}$ % | $ \Delta c_p/c_p _{\max}$ % | $ \Delta w/w _{\max}$ % |
|-----------------------------|--------------------|----------------------------|----------------------------|----------------------------|--------------------------------|----------------------------|
| $T_{3ab}(p)$ | 3a, 3b | 0.00075 | 0.00020 | 0.00020 | 0.0012 | 0.0010 |
| | 3d, 3e | 0.00061 | 0.00017 | 0.00013 | 0.0016 | 0.0005 |
| $T_{3cd}(p)$ | 3c, 3d | 0.00089 | 0.00027 | 0.00021 | 0.0040 | 0.0016 |
| | 3c, 3g | 0.00029 | 0.00009 | 0.00007 | 0.0017 | 0.0007 |
| | 3c, 3l | 0.00059 | 0.00019 | 0.00014 | 0.0039 | 0.0015 |
| | 3c, 3q | 0.00056 | 0.00018 | 0.00014 | 0.0040 | 0.0015 |
| | 3c, 3s | 0.00039 | 0.00012 | 0.00010 | 0.0031 | 0.0011 |
| $T_{3ef}(p)$ | 3e, 3f | 0.00060 | 0.00016 | 0.00012 | 0.0005 | 0.0001 |
| | 3h, 3i | 0.00061 | 0.00016 | 0.00012 | 0.0007 | 0.0001 |
| | 3n, 3o | 0.00031 | 0.00008 | 0.00006 | 0.0004 | 0.0001 |
| $T_{3gh}(p)$ | 3g, 3h | 0.00083 | 0.00022 | 0.00016 | 0.0058 | 0.0006 |
| | 3l, 3h | 0.00083 | 0.00022 | 0.00016 | 0.0064 | 0.0006 |
| | 3l, 3m | 0.00052 | 0.00014 | 0.00011 | 0.0058 | 0.0006 |
| $T_{3ij}(p)$ | 3i, 3j | 0.00034 | 0.00009 | 0.00007 | 0.0010 | 0.0002 |
| | 3p, 3j | 0.00036 | 0.00009 | 0.00007 | 0.0020 | 0.0002 |
| $T_{3jk}(p)$ | 3j, 3k | 0.00030 | 0.00007 | 0.00006 | 0.0008 | 0.0001 |
| | 3r, 3k | 0.00029 | 0.00007 | 0.00006 | 0.0018 | 0.0002 |
| $T_{3mn}(p)$ | 3m, 3n | 0.00090 | 0.00024 | 0.00017 | 0.0070 | 0.0003 |
| $T_{3op}(p)$ | 3o, 3p | 0.00041 | 0.00011 | 0.00008 | 0.0013 | 0.0002 |
| permissible tolerance | | 0.001 | 0.001 | 0.001 | 0.01 | 0.01 |

For example, the maximum relative difference between the backward equation of subregion 3a and the backward equation of subregion 3b along the subregion boundary $T_{3ab}(p)$ was determined as follows:

$$\left| \frac{\Delta v}{v} \right|_{\max} = \left| \frac{v_{3a}(p, T_{3ab}(p)) - v_{3b}(p, T_{3ab}(p))}{v_{3b}(p, T_{3ab}(p))} \right|_{\max}.$$

In addition, Tables 7 and 8 contain the maximum relative differences of specific enthalpy, specific entropy, specific isobaric heat capacity and speed of sound, calculated as described in Section 5.3, along the subregion boundaries of the $v(p, T)$ backward equations. For example, the maximum relative difference of specific enthalpy along the subregion boundary $T_{3ab}(p)$ was determined as follows:

$$\left| \frac{\Delta h}{h} \right|_{\max} = \left| \frac{h_3^{97}(v_{3a}, T_{3ab}) - h_3^{97}(v_{3b}, T_{3ab})}{h_3^{97}(v_{3b}, T_{3ab})} \right|_{\max}$$

where $v_{3a} = v_{3a}(p, T_{3ab}(p))$ and $v_{3b} = v_{3b}(p, T_{3ab}(p))$.

Tables 7 and 8 show that the relative specific volume differences between the backward equations $v(p,T)$ of the adjacent subregions and the maximum relative deviations of specific enthalpy, specific entropy, specific isobaric heat capacity, and speed of sound along the subregion boundary pressures and along the subregion boundary equations are smaller than the permissible numerical tolerances of these equations with the IAPWS-IF97 basic equation.

6 Auxiliary Equations $v(p,T)$ for the Region very close to the Critical Point

6.1 Subregions

The auxiliary equations $v(p,T)$ for the subregions 3u to 3z are valid from

$$T_{3qu}(p) < T \leq T_{3rx}(p) \text{ for } p_{\text{sat}}^{97}(643.15 \text{ K}) < p \leq 22.5 \text{ MPa}; \text{ see Figure 5.}$$




Figure 5. Division of region 3 into subregions 3u to 3z for the auxiliary equations

The subregion boundary equation $T_{3uv}(p)$ has the form of Eq. (1) and $T_{3wx}(p)$ has the form of Eq. (2). The coefficients n_i and the exponents I_i of the boundary equations are listed in Table 9.

Table 9. Numerical values of the coefficients of the equations $T_{3uv}(p)$ and $T_{3wx}(p)$ for subregion boundaries

| Equation | i | I_i | n_i | i | I_i | n_i |
|--------------|-----|-------|---|-----|-------|--|
| $T_{3uv}(p)$ | 1 | 0 | $0.528\ 199\ 646\ 263\ 062 \times 10^3$ | 3 | 2 | $-0.222\ 814\ 134\ 903\ 755$ |
| | 2 | 1 | $0.890\ 579\ 602\ 135\ 307 \times 10^1$ | 4 | 3 | $0.286\ 791\ 682\ 263\ 697 \times 10^{-2}$ |
| $T_{3wx}(p)$ | 1 | 0 | $0.728\ 052\ 609\ 145\ 380 \times 10^1$ | 4 | -1 | $0.329\ 196\ 213\ 998\ 375 \times 10^3$ |
| | 2 | 1 | $0.973\ 505\ 869\ 861\ 952 \times 10^2$ | 5 | -2 | $0.873\ 371\ 668\ 682\ 417 \times 10^3$ |
| | 3 | 2 | $0.147\ 370\ 491\ 183\ 191 \times 10^2$ | | | |

The following description of the use of the subregion boundary equations is summarized in Table 10 and Figure 5.

Table 10. Pressure ranges and corresponding subregion boundary equations for determining the correct subregion, 3u to 3z, for the auxiliary equations $v(p, T)$

| Supercritical Pressure Region | | | | |
|---|---|----------------------------------|------------|----------------------------------|
| Pressure Range | Sub-region | For | Sub-region | For |
| $22.11\ \text{MPa} < p \leq 22.5\ \text{MPa}$ | 3u | $T_{3qu}(p) < T \leq T_{3uv}(p)$ | 3v | $T_{3uv}(p) < T \leq T_{3ef}(p)$ |
| | 3w | $T_{3ef}(p) < T \leq T_{3wx}(p)$ | 3x | $T_{3wx}(p) < T \leq T_{3rx}(p)$ |
| $22.064\ \text{MPa} < p \leq 22.11\ \text{MPa}$ | 3u | $T_{3qu}(p) < T \leq T_{3uv}(p)$ | 3y | $T_{3uv}(p) < T \leq T_{3ef}(p)$ |
| | 3z | $T_{3ef}(p) < T \leq T_{3wx}(p)$ | 3x | $T_{3wx}(p) < T \leq T_{3rx}(p)$ |
| Subcritical Pressure Region | | | | |
| Temperature Range | Pressure Range | | Sub-region | For |
| $T \leq T_{\text{sat}}^{97}(p)$ | $p_{\text{sat}}^{97}(0.00264\ \text{m}^3\ \text{kg}^{-1})^e < p \leq 22.064\ \text{MPa}$ | | 3u | $T_{3qu}(p) < T \leq T_{3uv}(p)$ |
| | | | 3y | $T_{3uv}(p) < T$ |
| | $p_{\text{sat}}^{97}(643.15\ \text{K}) < p \leq p_{\text{sat}}^{97}(0.00264\ \text{m}^3\ \text{kg}^{-1})^e$ | | 3u | $T_{3qu}(p) < T$ |
| $T \geq T_{\text{sat}}^{97}(p)$ | $p_{\text{sat}}^{97}(0.00385\ \text{m}^3\ \text{kg}^{-1})^f < p \leq 22.064\ \text{MPa}$ | | 3z | $T \leq T_{3wx}(p)$ |
| | | | 3x | $T_{3wx}(p) < T \leq T_{3rx}(p)$ |
| | | | 3x | $T \leq T_{3rx}(p)$ |
| | $p_{\text{sat}}^{97}(643.15\ \text{K}) < p \leq p_{\text{sat}}^{97}(0.00385\ \text{m}^3\ \text{kg}^{-1})^f$ | | 3x | $T \leq T_{3rx}(p)$ |

$$^e p_{\text{sat}}^{97}(0.00264\ \text{m}^3\ \text{kg}^{-1}) = 2.193\ 161\ 551 \times 10^1\ \text{MPa}$$

$$^f p_{\text{sat}}^{97}(0.00385\ \text{m}^3\ \text{kg}^{-1}) = 2.190\ 096\ 265 \times 10^1\ \text{MPa}$$

The equation $T_{3uv}(p)$ approximates the isochore $v = 0.00264 \text{ m}^3 \text{ kg}^{-1}$ from $p = p_{\text{sat}}^{97}(0.00264 \text{ m}^3 \text{ kg}^{-1})$, where $p_{\text{sat}}^{97}(0.00264 \text{ m}^3 \text{ kg}^{-1}) = 2.193\,161\,551 \times 10^1 \text{ MPa}$, to 22.5 MPa and represents the boundary equation between subregions 3v or 3y and subregion 3u.

The equation $T_{3wx}(p)$ approximates the isochore $v = 0.00385 \text{ m}^3 \text{ kg}^{-1}$ from $p = p_{\text{sat}}^{97}(0.00385 \text{ m}^3 \text{ kg}^{-1})$, where $p_{\text{sat}}^{97}(0.00385 \text{ m}^3 \text{ kg}^{-1}) = 2.190\,096\,265 \times 10^1 \text{ MPa}$, to 22.5 MPa and represents the boundary equation between subregion 3x and subregions 3w or 3z.

Computer-program verification

To assist the user in computer-program verification of the equations for the subregion boundaries, Table 11 contains test values for calculated temperatures.

Table 11. Selected temperature values calculated from the subregion boundary equations $T_{3uv}(p)$ and $T_{3wx}(p)$ ^g

| Equation | p MPa | T K |
|--------------|------------|-------------------------------|
| $T_{3uv}(p)$ | 22.3 | $6.477\,996\,121 \times 10^2$ |
| $T_{3wx}(p)$ | 22.3 | $6.482\,049\,480 \times 10^2$ |

^g It is recommended that programmed functions be verified using 8 byte real values for all variables.

6.2 Auxiliary Equations $v(p,T)$ for the Subregions 3u to 3z

The auxiliary equations $v(p,T)$ for the subregions 3u to 3z have the dimensionless form of Eq. (4). The reducing quantities v^* , p^* , and T^* , the number of coefficients N , the non-linear parameters a and b , and the exponents c , d , and e are listed in Table 12 for the auxiliary equations of the subregions 3u to 3z. The coefficients n_i and exponents I_i and J_i are listed in Tables A2.1 to A2.6 of the Appendix.

Table 12. Reducing quantities v^* , p^* , and T^* , number of coefficients N , non-linear parameters a and b , and exponents c , d , and e for the auxiliary equations $v(p,T)$ of the subregions 3u to 3z

| Subregion | v^* $\text{m}^3 \text{ kg}^{-1}$ | p^* MPa | T^* K | N | a | b | c | d | e |
|-----------|---------------------------------------|--------------|------------|-----|-------|-------|-----|-----|-----|
| 3u | 0.0026 | 23 | 650 | 38 | 0.902 | 0.988 | 1 | 1 | 1 |
| 3v | 0.0031 | 23 | 650 | 39 | 0.960 | 0.995 | 1 | 1 | 1 |
| 3w | 0.0039 | 23 | 650 | 35 | 0.959 | 0.995 | 1 | 1 | 4 |
| 3x | 0.0049 | 23 | 650 | 36 | 0.910 | 0.988 | 1 | 1 | 1 |
| 3y | 0.0031 | 22 | 650 | 20 | 0.996 | 0.994 | 1 | 1 | 4 |
| 3z | 0.0038 | 22 | 650 | 23 | 0.993 | 0.994 | 1 | 1 | 4 |

Computer-program verification

To assist the user in computer-program verification of the auxiliary equations for the subregions 3u to 3z, Table 13 contains test values for calculated specific volumes.

Table 13. Selected specific volume values calculated from the auxiliary equations for the subregions 3u to 3z^h

| Equation | p MPa | T K | v $\text{m}^3 \text{kg}^{-1}$ | Equation | p MPa | T K | v $\text{m}^3 \text{kg}^{-1}$ |
|---------------|------------|----------|------------------------------------|---------------|------------|----------|------------------------------------|
| $v_{3u}(p,T)$ | 21.5 | 644.6 | $2.268\ 366\ 647 \times 10^{-3}$ | $v_{3x}(p,T)$ | 22.11 | 648.0 | $4.528\ 072\ 649 \times 10^{-3}$ |
| | 22.0 | 646.1 | $2.296\ 350\ 553 \times 10^{-3}$ | | 22.3 | 649.0 | $4.556\ 905\ 799 \times 10^{-3}$ |
| $v_{3v}(p,T)$ | 22.5 | 648.6 | $2.832\ 373\ 260 \times 10^{-3}$ | $v_{3y}(p,T)$ | 22.0 | 646.84 | $2.698\ 354\ 719 \times 10^{-3}$ |
| | 22.3 | 647.9 | $2.811\ 424\ 405 \times 10^{-3}$ | | 22.064 | 647.05 | $2.717\ 655\ 648 \times 10^{-3}$ |
| $v_{3w}(p,T)$ | 22.15 | 647.5 | $3.694\ 032\ 281 \times 10^{-3}$ | $v_{3z}(p,T)$ | 22.0 | 646.89 | $3.798\ 732\ 962 \times 10^{-3}$ |
| | 22.3 | 648.1 | $3.622\ 226\ 305 \times 10^{-3}$ | | 22.064 | 647.15 | $3.701\ 940\ 010 \times 10^{-3}$ |

^h It is recommended that programmed functions be verified using 8 byte real values for all variables.

6.3 Numerical Consistency

6.3.1 Numerical Consistency with the Basic Equation of IAPWS-IF97

The maximum relative differences and root-mean-square relative deviations of specific volume, calculated from the auxiliary equations $v(p,T)$ for subregions 3u to 3z, to the IAPWS-IF97 basic equation $f_3^{97}(v,T)$ are listed in Table 14. For the calculation of the root-mean-square values, which is described in Section 1, one million points uniformly distributed over the range of validity in the p - T plane have been used.

Table 14 shows that the deviations of the specific volume from the IAPWS-IF97 basic equation are better than 0.1 %. Only in a small region for pressures less than 22.11 MPa (see Figure 5) do the deviations of the specific volume from the IAPWS-IF97 basic equation approach 2 %.

Table 14. Maximum relative deviations and root-mean-square relative deviations of the specific volume, calculated from the auxiliary equations for subregions 3u to 3z from the IAPWS-IF97 basic equation

| Subregion | $ \Delta v/v $ % | | Subregion | $ \Delta v/v $ % | |
|-----------|---------------------|-------|-----------|---------------------|-------|
| | max | RMS | | max | RMS |
| 3u | 0.097 | 0.058 | 3x | 0.090 | 0.050 |
| 3v | 0.082 | 0.040 | 3y | 1.77 | 1.04 |
| 3w | 0.065 | 0.023 | 3z | 1.80 | 0.921 |

6.3.2 Consistency at Boundaries Between Subregions

The maximum relative differences of specific volume between the $v(p,T)$ auxiliary equations of adjacent subregions along the subregion boundary pressures are listed in Table 15. Table 16 contains these maximum relative differences along the subregion boundary equations.

Table 15. Maximum relative deviations of specific volume between the auxiliary equations $v(p,T)$ of the adjacent subregions along the subregion boundary pressures

| Subregion Boundary | Between Subregions | $ \Delta v/v _{\max}$ % |
|--------------------|--------------------|----------------------------|
| $p = 22.5$ MPa | 3l, 3u | 0.096 |
| | 3m, 3u | 0.096 |
| | 3m, 3v | 0.035 |
| | 3n, 3v | 0.046 |
| | 3o, 3w | 0.019 |
| | 3p, 3w | 0.021 |
| | 3p, 3x | 0.042 |
| | 3j, 3x | 0.043 |
| $p = 22.11$ MPa | 3v, 3y | 1.7 |
| | 3w, 3z | 1.7 |

Table 16. Maximum relative deviations of specific volume between the auxiliary equations $v(p,T)$ of the adjacent subregions along the subregion boundary equations

| Subregion Boundary Equation | Between Subregions | $ \Delta v/v _{\max}$ % |
|-----------------------------|--------------------|----------------------------|
| $T_{3qu}(p)$ | 3q, 3u | 0.097 |
| $T_{3rx}(p)$ | 3x, 3r | 0.045 |
| $T_{3uv}(p)$ | 3u, 3v | 0.14 |
| | 3u, 3y | 1.8 |
| $T_{3ef}(p)$ | 3v, 3w | 0.080 |
| | 3y, 3z | 3.5 |
| $T_{3wx}(p)$ | 3w, 3x | 0.049 |
| | 3z, 3x | 1.8 |

7 Computing Time in Relation to IAPWS-IF97

A very important motivation for the development of the backward equations $v(p,T)$ was reducing the computing time to obtain thermodynamic properties and differential quotients from given variables (p,T) in region 3. Using IAPWS-IF97, time-consuming iteration is required. Using the $v(p,T)$ backward equations, iteration can be avoided. The calculation speed is about 17 times faster than iteration with IAPWS-IF97.

If iteration is used, the time to reach convergence can be significantly reduced by using the backward equations $v(p,T)$ to calculate very accurate starting values.

8 Application of the Backward and Auxiliary Equations $v(p,T)$

The numerical consistency of the specific volume v calculated from the main backward equations $v_3(p,T)$ described in Section 5 with the IAPWS-IF97 basic equation $f_3^{97}(v,T)$ is sufficient for most applications in process modeling. For many calculations, the numerical consistency of the auxiliary equations described in Section 6 is also satisfactory in the region very close to the critical point.

For applications where the demands on numerical consistency are extremely high, iteration using the IAPWS-IF97 basic equation $f(v,T)$ may be necessary. In these cases, the backward and auxiliary equations $v(p,T)$ can be used for calculating very accurate starting values.

The backward and auxiliary equations $v(p,T)$ should only be used in their ranges of validity described in Section 4. They should not be used for determining any thermodynamic derivatives. They should also not be used together with the fundamental equation in iterative calculations of other backward functions such as $T(p,h)$ or $T(p,s)$. Iteration of backward functions can only be performed by using the fundamental equations.

In any case, depending on the application, a conscious decision is required whether to use the backward and auxiliary equations $v(p,T)$ or to calculate the corresponding values by iteration from the basic equation of IAPWS-IF97.

9 References

- [1] IAPWS, *Revised Release on the IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam* (2007), available from: <http://www.iapws.org>.
- [2] Wagner, W., Cooper, J. R., Dittmann, A., Kijima, J., Kretschmar, H.-J., Kruse, A., Mareš, R., Oguchi, K., Sato, H., Stöcker, I., Šifner, O., Tanishita, I., Trübenbach, J., and Willkommen, Th., *The IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam*, *ASME J. Eng. Gas Turbines Power* **122**, 150-182 (2000).
- [3] Kretschmar, H.-J., Harvey, A. H., Knobloch, K., Mareš, R., Miyagawa, K., Okita, N., Span, R., Stöcker, I., Wagner, W., and Weber, I., *Supplementary Backward Equations $v(p,T)$ for the Critical and Supercritical Regions (Region 3) of the IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam*, *ASME J. Eng. Gas Turbines Power* **131**, 043101 (2009).
- [4] IAPWS, *Revised Supplementary Release on Backward Equations for Pressure as a Function of Enthalpy and Entropy $p(h,s)$ for Regions 1 and 2 of the IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam* (2014), available from <http://www.iapws.org>.

- [5] Kretzschmar, H.-J., Cooper, J. R., Dittmann, A., Friend, D. G., Gallagher, J. S., Knobloch, K., Mareš, R., Miyagawa, K., Stöcker, I., Trübenbach, J., Wagner, W., and Willkommen, Th., Supplementary Backward Equations for Pressure as a Function of Enthalpy and Entropy $p(h,s)$ to the Industrial Formulation IAPWS-IF97 for Water and Steam, *ASME J. Eng. Gas Turbines Power* **128**, 702-713 (2006).
- [6] IAPWS, *Revised Supplementary Release on Backward Equations for the Functions $T(p,h)$, $v(p,h)$ and $T(p,s)$, $v(p,s)$ for Region 3 of the IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam* (2014), available from: <http://www.iapws.org>.
- [7] Kretzschmar, H.-J., Cooper, J. R., Dittmann, A., Friend, D. G., Gallagher, J. S., Harvey, A. H., Knobloch, K., Mareš, R., Miyagawa, K., Okita, N., Stöcker, I., Wagner, W., and Weber, I., Supplementary Backward Equations $T(p,h)$, $v(p,h)$, and $T(p,s)$, $v(p,s)$ for the Critical and Supercritical Regions (Region 3) of the Industrial Formulation IAPWS-IF97 for Water and Steam, *ASME J. Eng. Gas Turbines Power* **129**, 294-303 (2007).
- [8] IAPWS, *Revised Supplementary Release on Backward Equations $p(h,s)$ for Region 3, Equations as a Function of h and s for the Region Boundaries, and an Equation $T_{sat}(h,s)$ for Region 4 of the IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam* (2014), available from: <http://www.iapws.org>.
- [9] Kretzschmar, H.-J., Cooper, J. R., Dittmann, A., Friend, D. G., Gallagher, J. S., Harvey, A. H., Knobloch, K., Mareš, R., Miyagawa, K., Okita, N., Span, R., Stöcker, I., Wagner, W., and Weber, I., Supplementary Backward Equations $p(h, s)$ for the Critical and Supercritical Regions (Region 3), and Equations for the Two-Phase Region and Region Boundaries of the IAPWS Industrial Formulation 1997 for the Thermodynamic Properties of Water and Steam, *ASME J. Eng. Gas Turbines Power* **129**, 1125-1137 (2007).
- [10] IAPWS, *Release on the Values of Temperature, Pressure and Density of Ordinary and Heavy Water Substances at their Respective Critical Points* (1992), available from <http://www.iapws.org>.

Appendix

A1 Coefficients for Backward Equations

Table A1.1. Coefficients and exponents of the backward equation $v_{3a}(p, T)$ for subregion 3a

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|---|
| 1 | -12 | 5 | $0.110\ 879\ 558\ 823\ 853 \times 10^{-2}$ | 16 | -3 | 1 | $-0.122\ 494\ 831\ 387\ 441 \times 10^{-1}$ |
| 2 | -12 | 10 | $0.572\ 616\ 740\ 810\ 616 \times 10^3$ | 17 | -3 | 3 | $0.179\ 357\ 604\ 019\ 989 \times 10^1$ |
| 3 | -12 | 12 | $-0.767\ 051\ 948\ 380\ 852 \times 10^5$ | 18 | -3 | 6 | $0.442\ 729\ 521\ 058\ 314 \times 10^2$ |
| 4 | -10 | 5 | $-0.253\ 321\ 069\ 529\ 674 \times 10^{-1}$ | 19 | -2 | 0 | $-0.593\ 223\ 489\ 018\ 342 \times 10^{-2}$ |
| 5 | -10 | 10 | $0.628\ 008\ 049\ 345\ 689 \times 10^4$ | 20 | -2 | 2 | $0.453\ 186\ 261\ 685\ 774$ |
| 6 | -10 | 12 | $0.234\ 105\ 654\ 131\ 876 \times 10^6$ | 21 | -2 | 3 | $0.135\ 825\ 703\ 129\ 140 \times 10^1$ |
| 7 | -8 | 5 | $0.216\ 867\ 826\ 045\ 856$ | 22 | -1 | 0 | $0.408\ 748\ 415\ 856\ 745 \times 10^{-1}$ |
| 8 | -8 | 8 | $-0.156\ 237\ 904\ 341\ 963 \times 10^3$ | 23 | -1 | 1 | $0.474\ 686\ 397\ 863\ 312$ |
| 9 | -8 | 10 | $-0.269\ 893\ 956\ 176\ 613 \times 10^5$ | 24 | -1 | 2 | $0.118\ 646\ 814\ 997\ 915 \times 10^1$ |
| 10 | -6 | 1 | $-0.180\ 407\ 100\ 085\ 505 \times 10^{-3}$ | 25 | 0 | 0 | $0.546\ 987\ 265\ 727\ 549$ |
| 11 | -5 | 1 | $0.116\ 732\ 227\ 668\ 261 \times 10^{-2}$ | 26 | 0 | 1 | $0.195\ 266\ 770\ 452\ 643$ |
| 12 | -5 | 5 | $0.266\ 987\ 040\ 856\ 040 \times 10^2$ | 27 | 1 | 0 | $-0.502\ 268\ 790\ 869\ 663 \times 10^{-1}$ |
| 13 | -5 | 10 | $0.282\ 776\ 617\ 243\ 286 \times 10^5$ | 28 | 1 | 2 | $-0.369\ 645\ 308\ 193\ 377$ |
| 14 | -4 | 8 | $-0.242\ 431\ 520\ 029\ 523 \times 10^4$ | 29 | 2 | 0 | $0.633\ 828\ 037\ 528\ 420 \times 10^{-2}$ |
| 15 | -3 | 0 | $0.435\ 217\ 323\ 022\ 733 \times 10^{-3}$ | 30 | 2 | 2 | $0.797\ 441\ 793\ 901\ 017 \times 10^{-1}$ |

Table A1.2. Coefficients and exponents of the backward equation $v_{3b}(p, T)$ for subregion 3b

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|---|
| 1 | -12 | 10 | $-0.827\ 670\ 470\ 003\ 621 \times 10^{-1}$ | 17 | -3 | 2 | $-0.416\ 375\ 290\ 166\ 236 \times 10^{-1}$ |
| 2 | -12 | 12 | $0.416\ 887\ 126\ 010\ 565 \times 10^2$ | 18 | -3 | 3 | $-0.413\ 754\ 957\ 011\ 042 \times 10^2$ |
| 3 | -10 | 8 | $0.483\ 651\ 982\ 197\ 059 \times 10^{-1}$ | 19 | -3 | 5 | $-0.506\ 673\ 295\ 721\ 637 \times 10^2$ |
| 4 | -10 | 14 | $-0.291\ 032\ 084\ 950\ 276 \times 10^5$ | 20 | -2 | 0 | $-0.572\ 212\ 965\ 569\ 023 \times 10^{-3}$ |
| 5 | -8 | 8 | $-0.111\ 422\ 582\ 236\ 948 \times 10^3$ | 21 | -2 | 2 | $0.608\ 817\ 368\ 401\ 785 \times 10^1$ |
| 6 | -6 | 5 | $-0.202\ 300\ 083\ 904\ 014 \times 10^{-1}$ | 22 | -2 | 5 | $0.239\ 600\ 660\ 256\ 161 \times 10^2$ |
| 7 | -6 | 6 | $0.294\ 002\ 509\ 338\ 515 \times 10^3$ | 23 | -1 | 0 | $0.122\ 261\ 479\ 925\ 384 \times 10^{-1}$ |
| 8 | -6 | 8 | $0.140\ 244\ 997\ 609\ 658 \times 10^3$ | 24 | -1 | 2 | $0.216\ 356\ 057\ 692\ 938 \times 10^1$ |
| 9 | -5 | 5 | $-0.344\ 384\ 158\ 811\ 459 \times 10^3$ | 25 | 0 | 0 | $0.398\ 198\ 903\ 368\ 642$ |
| 10 | -5 | 8 | $0.361\ 182\ 452\ 612\ 149 \times 10^3$ | 26 | 0 | 1 | $-0.116\ 892\ 827\ 834\ 085$ |
| 11 | -5 | 10 | $-0.140\ 699\ 677\ 420\ 738 \times 10^4$ | 27 | 1 | 0 | $-0.102\ 845\ 919\ 373\ 532$ |
| 12 | -4 | 2 | $-0.202\ 023\ 902\ 676\ 481 \times 10^{-2}$ | 28 | 1 | 2 | $-0.492\ 676\ 637\ 589\ 284$ |
| 13 | -4 | 4 | $0.171\ 346\ 792\ 457\ 471 \times 10^3$ | 29 | 2 | 0 | $0.655\ 540\ 456\ 406\ 790 \times 10^{-1}$ |
| 14 | -4 | 5 | $-0.425\ 597\ 804\ 058\ 632 \times 10^1$ | 30 | 3 | 2 | $-0.240\ 462\ 535\ 078\ 530$ |
| 15 | -3 | 0 | $0.691\ 346\ 085\ 000\ 334 \times 10^{-5}$ | 31 | 4 | 0 | $-0.269\ 798\ 180\ 310\ 075 \times 10^{-1}$ |
| 16 | -3 | 1 | $0.151\ 140\ 509\ 678\ 925 \times 10^{-2}$ | 32 | 4 | 1 | $0.128\ 369\ 435\ 967\ 012$ |

Table A1.3. Coefficients and exponents of the backward equation $v_{3c}(p, T)$ for subregion 3c

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|---|
| 1 | -12 | 6 | 0.311 967 788 763 030 $\times 10^1$ | 19 | -2 | 4 | 0.234 604 891 591 616 $\times 10^3$ |
| 2 | -12 | 8 | 0.276 713 458 847 564 $\times 10^5$ | 20 | -2 | 5 | 0.377 515 668 966 951 $\times 10^4$ |
| 3 | -12 | 10 | 0.322 583 103 403 269 $\times 10^8$ | 21 | -1 | 0 | 0.158 646 812 591 361 $\times 10^{-1}$ |
| 4 | -10 | 6 | -0.342 416 065 095 363 $\times 10^3$ | 22 | -1 | 1 | 0.707 906 336 241 843 |
| 5 | -10 | 8 | -0.899 732 529 907 377 $\times 10^6$ | 23 | -1 | 2 | 0.126 016 225 146 570 $\times 10^2$ |
| 6 | -10 | 10 | -0.793 892 049 821 251 $\times 10^8$ | 24 | 0 | 0 | 0.736 143 655 772 152 |
| 7 | -8 | 5 | 0.953 193 003 217 388 $\times 10^2$ | 25 | 0 | 1 | 0.676 544 268 999 101 |
| 8 | -8 | 6 | 0.229 784 742 345 072 $\times 10^4$ | 26 | 0 | 2 | -0.178 100 588 189 137 $\times 10^2$ |
| 9 | -8 | 7 | 0.175 336 675 322 499 $\times 10^6$ | 27 | 1 | 0 | -0.156 531 975 531 713 |
| 10 | -6 | 8 | 0.791 214 365 222 792 $\times 10^7$ | 28 | 1 | 2 | 0.117 707 430 048 158 $\times 10^2$ |
| 11 | -5 | 1 | 0.319 933 345 844 209 $\times 10^{-4}$ | 29 | 2 | 0 | 0.840 143 653 860 447 $\times 10^{-1}$ |
| 12 | -5 | 4 | -0.659 508 863 555 767 $\times 10^2$ | 30 | 2 | 1 | -0.186 442 467 471 949 |
| 13 | -5 | 7 | -0.833 426 563 212 851 $\times 10^6$ | 31 | 2 | 3 | -0.440 170 203 949 645 $\times 10^2$ |
| 14 | -4 | 2 | 0.645 734 680 583 292 $\times 10^{-1}$ | 32 | 2 | 7 | 0.123 290 423 502 494 $\times 10^7$ |
| 15 | -4 | 8 | -0.382 031 020 570 813 $\times 10^7$ | 33 | 3 | 0 | -0.240 650 039 730 845 $\times 10^{-1}$ |
| 16 | -3 | 0 | 0.406 398 848 470 079 $\times 10^{-4}$ | 34 | 3 | 7 | -0.107 077 716 660 869 $\times 10^7$ |
| 17 | -3 | 3 | 0.310 327 498 492 008 $\times 10^2$ | 35 | 8 | 1 | 0.438 319 858 566 475 $\times 10^{-1}$ |
| 18 | -2 | 0 | -0.892 996 718 483 724 $\times 10^{-3}$ | | | | |

Table A1.4. Coefficients and exponents of the backward equation $v_{3d}(p, T)$ for subregion 3d

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|--|-----|-------|-------|---|
| 1 | -12 | 4 | -0.452 484 847 171 645 $\times 10^{-9}$ | 20 | -5 | 1 | -0.436 701 347 922 356 $\times 10^{-5}$ |
| 2 | -12 | 6 | 0.315 210 389 538 801 $\times 10^{-4}$ | 21 | -5 | 2 | -0.404 213 852 833 996 $\times 10^{-3}$ |
| 3 | -12 | 7 | -0.214 991 352 047 545 $\times 10^{-2}$ | 22 | -5 | 5 | -0.348 153 203 414 663 $\times 10^3$ |
| 4 | -12 | 10 | 0.508 058 874 808 345 $\times 10^3$ | 23 | -5 | 7 | -0.385 294 213 555 289 $\times 10^6$ |
| 5 | -12 | 12 | -0.127 123 036 845 932 $\times 10^8$ | 24 | -4 | 0 | 0.135 203 700 099 403 $\times 10^{-6}$ |
| 6 | -12 | 16 | 0.115 371 133 120 497 $\times 10^{13}$ | 25 | -4 | 1 | 0.134 648 383 271 089 $\times 10^{-3}$ |
| 7 | -10 | 0 | -0.197 805 728 776 273 $\times 10^{-15}$ | 26 | -4 | 7 | 0.125 031 835 351 736 $\times 10^6$ |
| 8 | -10 | 2 | 0.241 554 806 033 972 $\times 10^{-10}$ | 27 | -3 | 2 | 0.968 123 678 455 841 $\times 10^{-1}$ |
| 9 | -10 | 4 | -0.156 481 703 640 525 $\times 10^{-5}$ | 28 | -3 | 4 | 0.225 660 517 512 438 $\times 10^3$ |
| 10 | -10 | 6 | 0.277 211 346 836 625 $\times 10^{-2}$ | 29 | -2 | 0 | -0.190 102 435 341 872 $\times 10^{-3}$ |
| 11 | -10 | 8 | -0.203 578 994 462 286 $\times 10^2$ | 30 | -2 | 1 | -0.299 628 410 819 229 $\times 10^{-1}$ |
| 12 | -10 | 10 | 0.144 369 489 909 053 $\times 10^7$ | 31 | -1 | 0 | 0.500 833 915 372 121 $\times 10^{-2}$ |
| 13 | -10 | 14 | -0.411 254 217 946 539 $\times 10^{11}$ | 32 | -1 | 1 | 0.387 842 482 998 411 |
| 14 | -8 | 3 | 0.623 449 786 243 773 $\times 10^{-5}$ | 33 | -1 | 5 | -0.138 535 367 777 182 $\times 10^4$ |
| 15 | -8 | 7 | -0.221 774 281 146 038 $\times 10^2$ | 34 | 0 | 0 | 0.870 745 245 971 773 |
| 16 | -8 | 8 | -0.689 315 087 933 158 $\times 10^5$ | 35 | 0 | 2 | 0.171 946 252 068 742 $\times 10^1$ |
| 17 | -8 | 10 | -0.195 419 525 060 713 $\times 10^8$ | 36 | 1 | 0 | -0.326 650 121 426 383 $\times 10^{-1}$ |
| 18 | -6 | 6 | 0.316 373 510 564 015 $\times 10^4$ | 37 | 1 | 6 | 0.498 044 171 727 877 $\times 10^4$ |
| 19 | -6 | 8 | 0.224 040 754 426 988 $\times 10^7$ | 38 | 3 | 0 | 0.551 478 022 765 087 $\times 10^{-2}$ |

Table A1.5. Coefficients and exponents of the backward equation $v_{3e}(p, T)$ for subregion 3e

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|---|
| 1 | -12 | 14 | $0.715\ 815\ 808\ 404\ 721 \times 10^9$ | 16 | -3 | 6 | $0.475\ 992\ 667\ 717\ 124 \times 10^5$ |
| 2 | -12 | 16 | $-0.114\ 328\ 360\ 753\ 449 \times 10^{12}$ | 17 | -3 | 7 | $-0.266\ 627\ 750\ 390\ 341 \times 10^6$ |
| 3 | -10 | 3 | $0.376\ 531\ 002\ 015\ 720 \times 10^{-11}$ | 18 | -2 | 0 | $-0.153\ 314\ 954\ 386\ 524 \times 10^{-3}$ |
| 4 | -10 | 6 | $-0.903\ 983\ 668\ 691\ 157 \times 10^{-4}$ | 19 | -2 | 1 | $0.305\ 638\ 404\ 828\ 265$ |
| 5 | -10 | 10 | $0.665\ 695\ 908\ 836\ 252 \times 10^6$ | 20 | -2 | 3 | $0.123\ 654\ 999\ 499\ 486 \times 10^3$ |
| 6 | -10 | 14 | $0.535\ 364\ 174\ 960\ 127 \times 10^{10}$ | 21 | -2 | 4 | $-0.104\ 390\ 794\ 213\ 011 \times 10^4$ |
| 7 | -10 | 16 | $0.794\ 977\ 402\ 335\ 603 \times 10^{11}$ | 22 | -1 | 0 | $-0.157\ 496\ 516\ 174\ 308 \times 10^{-1}$ |
| 8 | -8 | 7 | $0.922\ 230\ 563\ 421\ 437 \times 10^2$ | 23 | 0 | 0 | $0.685\ 331\ 118\ 940\ 253$ |
| 9 | -8 | 8 | $-0.142\ 586\ 073\ 991\ 215 \times 10^6$ | 24 | 0 | 1 | $0.178\ 373\ 462\ 873\ 903 \times 10^1$ |
| 10 | -8 | 10 | $-0.111\ 796\ 381\ 424\ 162 \times 10^7$ | 25 | 1 | 0 | $-0.544\ 674\ 124\ 878\ 910$ |
| 11 | -6 | 6 | $0.896\ 121\ 629\ 640\ 760 \times 10^4$ | 26 | 1 | 4 | $0.204\ 529\ 931\ 318\ 843 \times 10^4$ |
| 12 | -5 | 6 | $-0.669\ 989\ 239\ 070\ 491 \times 10^4$ | 27 | 1 | 6 | $-0.228\ 342\ 359\ 328\ 752 \times 10^5$ |
| 13 | -4 | 2 | $0.451\ 242\ 538\ 486\ 834 \times 10^{-2}$ | 28 | 2 | 0 | $0.413\ 197\ 481\ 515\ 899$ |
| 14 | -4 | 4 | $-0.339\ 731\ 325\ 977\ 713 \times 10^2$ | 29 | 2 | 2 | $-0.341\ 931\ 835\ 910\ 405 \times 10^2$ |
| 15 | -3 | 2 | $-0.120\ 523\ 111\ 552\ 278 \times 10^1$ | | | | |

Table A1.6. Coefficients and exponents of the backward equation $v_{3f}(p, T)$ for subregion 3f

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|--|
| 1 | 0 | -3 | $-0.251\ 756\ 547\ 792\ 325 \times 10^{-7}$ | 22 | 10 | -6 | $0.470\ 942\ 606\ 221\ 652 \times 10^{-5}$ |
| 2 | 0 | -2 | $0.601\ 307\ 193\ 668\ 763 \times 10^{-5}$ | 23 | 12 | -10 | $0.195\ 049\ 710\ 391\ 712 \times 10^{-12}$ |
| 3 | 0 | -1 | $-0.100\ 615\ 977\ 450\ 049 \times 10^{-2}$ | 24 | 12 | -8 | $-0.911\ 627\ 886\ 266\ 077 \times 10^{-8}$ |
| 4 | 0 | 0 | $0.999\ 969\ 140\ 252\ 192$ | 25 | 12 | -4 | $0.604\ 374\ 640\ 201\ 265 \times 10^{-3}$ |
| 5 | 0 | 1 | $0.214\ 107\ 759\ 236\ 486 \times 10^1$ | 26 | 14 | -12 | $-0.225\ 132\ 933\ 900\ 136 \times 10^{-15}$ |
| 6 | 0 | 2 | $-0.165\ 175\ 571\ 959\ 086 \times 10^2$ | 27 | 14 | -10 | $0.610\ 916\ 973\ 582\ 981 \times 10^{-11}$ |
| 7 | 1 | -1 | $-0.141\ 987\ 303\ 638\ 727 \times 10^{-2}$ | 28 | 14 | -8 | $-0.303\ 063\ 908\ 043\ 404 \times 10^{-6}$ |
| 8 | 1 | 1 | $0.269\ 251\ 915\ 156\ 554 \times 10^1$ | 29 | 14 | -6 | $-0.137\ 796\ 070\ 798\ 409 \times 10^{-4}$ |
| 9 | 1 | 2 | $0.349\ 741\ 815\ 858\ 722 \times 10^2$ | 30 | 14 | -4 | $-0.919\ 296\ 736\ 666\ 106 \times 10^{-3}$ |
| 10 | 1 | 3 | $-0.300\ 208\ 695\ 771\ 783 \times 10^2$ | 31 | 16 | -10 | $0.639\ 288\ 223\ 132\ 545 \times 10^{-9}$ |
| 11 | 2 | 0 | $-0.131\ 546\ 288\ 252\ 539 \times 10^1$ | 32 | 16 | -8 | $0.753\ 259\ 479\ 898\ 699 \times 10^{-6}$ |
| 12 | 2 | 1 | $-0.839\ 091\ 277\ 286\ 169 \times 10^1$ | 33 | 18 | -12 | $-0.400\ 321\ 478\ 682\ 929 \times 10^{-12}$ |
| 13 | 3 | -5 | $0.181\ 545\ 608\ 337\ 015 \times 10^{-9}$ | 34 | 18 | -10 | $0.756\ 140\ 294\ 351\ 614 \times 10^{-8}$ |
| 14 | 3 | -2 | $-0.591\ 099\ 206\ 478\ 909 \times 10^{-3}$ | 35 | 20 | -12 | $-0.912\ 082\ 054\ 034\ 891 \times 10^{-11}$ |
| 15 | 3 | 0 | $0.152\ 115\ 067\ 087\ 106 \times 10^1$ | 36 | 20 | -10 | $-0.237\ 612\ 381\ 140\ 539 \times 10^{-7}$ |
| 16 | 4 | -3 | $0.252\ 956\ 470\ 663\ 225 \times 10^{-4}$ | 37 | 20 | -6 | $0.269\ 586\ 010\ 591\ 874 \times 10^{-4}$ |
| 17 | 5 | -8 | $0.100\ 726\ 265\ 203\ 786 \times 10^{-14}$ | 38 | 22 | -12 | $-0.732\ 828\ 135\ 157\ 839 \times 10^{-10}$ |
| 18 | 5 | 1 | $-0.149\ 774\ 533\ 860\ 650 \times 10^1$ | 39 | 24 | -12 | $0.241\ 995\ 578\ 306\ 660 \times 10^{-9}$ |
| 19 | 6 | -6 | $-0.793\ 940\ 970\ 562\ 969 \times 10^{-9}$ | 40 | 24 | -4 | $-0.405\ 735\ 532\ 730\ 322 \times 10^{-3}$ |
| 20 | 7 | -4 | $-0.150\ 290\ 891\ 264\ 717 \times 10^{-3}$ | 41 | 28 | -12 | $0.189\ 424\ 143\ 498\ 011 \times 10^{-9}$ |
| 21 | 7 | 1 | $0.151\ 205\ 531\ 275\ 133 \times 10^1$ | 42 | 32 | -12 | $-0.486\ 632\ 965\ 074\ 563 \times 10^{-9}$ |

Table A1.7. Coefficients and exponents of the backward equation $v_{3g}(p, T)$ for subregion 3g

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|---|
| 1 | -12 | 7 | $0.412\ 209\ 020\ 652\ 996 \times 10^{-4}$ | 20 | -2 | 3 | $-0.910\ 782\ 540\ 134\ 681 \times 10^2$ |
| 2 | -12 | 12 | $-0.114\ 987\ 238\ 280\ 587 \times 10^7$ | 21 | -2 | 5 | $0.135\ 033\ 227\ 281\ 565 \times 10^6$ |
| 3 | -12 | 14 | $0.948\ 180\ 885\ 032\ 080 \times 10^{10}$ | 22 | -2 | 14 | $-0.712\ 949\ 383\ 408\ 211 \times 10^{19}$ |
| 4 | -12 | 18 | $-0.195\ 788\ 865\ 718\ 971 \times 10^{18}$ | 23 | -2 | 24 | $-0.104\ 578\ 785\ 289\ 542 \times 10^{37}$ |
| 5 | -12 | 22 | $0.496\ 250\ 704\ 871\ 300 \times 10^{25}$ | 24 | -1 | 2 | $0.304\ 331\ 584\ 444\ 093 \times 10^2$ |
| 6 | -12 | 24 | $-0.105\ 549\ 884\ 548\ 496 \times 10^{29}$ | 25 | -1 | 8 | $0.593\ 250\ 797\ 959\ 445 \times 10^{10}$ |
| 7 | -10 | 14 | $-0.758\ 642\ 165\ 988\ 278 \times 10^{12}$ | 26 | -1 | 18 | $-0.364\ 174\ 062\ 110\ 798 \times 10^{28}$ |
| 8 | -10 | 20 | $-0.922\ 172\ 769\ 596\ 101 \times 10^{23}$ | 27 | 0 | 0 | $0.921\ 791\ 403\ 532\ 461$ |
| 9 | -10 | 24 | $0.725\ 379\ 072\ 059\ 348 \times 10^{30}$ | 28 | 0 | 1 | $-0.337\ 693\ 609\ 657\ 471$ |
| 10 | -8 | 7 | $-0.617\ 718\ 249\ 205\ 859 \times 10^2$ | 29 | 0 | 2 | $-0.724\ 644\ 143\ 758\ 508 \times 10^2$ |
| 11 | -8 | 8 | $0.107\ 555\ 033\ 344\ 858 \times 10^5$ | 30 | 1 | 0 | $-0.110\ 480\ 239\ 272\ 601$ |
| 12 | -8 | 10 | $-0.379\ 545\ 802\ 336\ 487 \times 10^8$ | 31 | 1 | 1 | $0.536\ 516\ 031\ 875\ 059 \times 10^1$ |
| 13 | -8 | 12 | $0.228\ 646\ 846\ 221\ 831 \times 10^{12}$ | 32 | 1 | 3 | $-0.291\ 441\ 872\ 156\ 205 \times 10^4$ |
| 14 | -6 | 8 | $-0.499\ 741\ 093\ 010\ 619 \times 10^7$ | 33 | 3 | 24 | $0.616\ 338\ 176\ 535\ 305 \times 10^{40}$ |
| 15 | -6 | 22 | $-0.280\ 214\ 310\ 054\ 101 \times 10^{31}$ | 34 | 5 | 22 | $-0.120\ 889\ 175\ 861\ 180 \times 10^{39}$ |
| 16 | -5 | 7 | $0.104\ 915\ 406\ 769\ 586 \times 10^7$ | 35 | 6 | 12 | $0.818\ 396\ 024\ 524\ 612 \times 10^{23}$ |
| 17 | -5 | 20 | $0.613\ 754\ 229\ 168\ 619 \times 10^{28}$ | 36 | 8 | 3 | $0.940\ 781\ 944\ 835\ 829 \times 10^9$ |
| 18 | -4 | 22 | $0.802\ 056\ 715\ 528\ 378 \times 10^{32}$ | 37 | 10 | 0 | $-0.367\ 279\ 669\ 545\ 448 \times 10^5$ |
| 19 | -3 | 7 | $-0.298\ 617\ 819\ 828\ 065 \times 10^8$ | 38 | 10 | 6 | $-0.837\ 513\ 931\ 798\ 655 \times 10^{16}$ |

Table A1.8. Coefficients and exponents of the backward equation $v_{3h}(p, T)$ for subregion 3h

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|--|-----|-------|-------|---|
| 1 | -12 | 8 | $0.561\ 379\ 678\ 887\ 577 \times 10^{-1}$ | 16 | -6 | 8 | $-0.656\ 174\ 421\ 999\ 594 \times 10^7$ |
| 2 | -12 | 12 | $0.774\ 135\ 421\ 587\ 083 \times 10^{10}$ | 17 | -5 | 2 | $0.156\ 362\ 212\ 977\ 396 \times 10^{-4}$ |
| 3 | -10 | 4 | $0.111\ 482\ 975\ 877\ 938 \times 10^{-8}$ | 18 | -5 | 3 | $-0.212\ 946\ 257\ 021\ 400 \times 10^1$ |
| 4 | -10 | 6 | $-0.143\ 987\ 128\ 208\ 183 \times 10^{-2}$ | 19 | -5 | 4 | $0.135\ 249\ 306\ 374\ 858 \times 10^2$ |
| 5 | -10 | 8 | $0.193\ 696\ 558\ 764\ 920 \times 10^4$ | 20 | -4 | 2 | $0.177\ 189\ 164\ 145\ 813$ |
| 6 | -10 | 10 | $-0.605\ 971\ 823\ 585\ 005 \times 10^9$ | 21 | -4 | 4 | $0.139\ 499\ 167\ 345\ 464 \times 10^4$ |
| 7 | -10 | 14 | $0.171\ 951\ 568\ 124\ 337 \times 10^{14}$ | 22 | -3 | 1 | $-0.703\ 670\ 932\ 036\ 388 \times 10^{-2}$ |
| 8 | -10 | 16 | $-0.185\ 461\ 154\ 985\ 145 \times 10^{17}$ | 23 | -3 | 2 | $-0.152\ 011\ 044\ 389\ 648$ |
| 9 | -8 | 0 | $0.387\ 851\ 168\ 078\ 010 \times 10^{-16}$ | 24 | -2 | 0 | $0.981\ 916\ 922\ 991\ 113 \times 10^{-4}$ |
| 10 | -8 | 1 | $-0.395\ 464\ 327\ 846\ 105 \times 10^{-13}$ | 25 | -1 | 0 | $0.147\ 199\ 658\ 618\ 076 \times 10^{-2}$ |
| 11 | -8 | 6 | $-0.170\ 875\ 935\ 679\ 023 \times 10^3$ | 26 | -1 | 2 | $0.202\ 618\ 487\ 025\ 578 \times 10^2$ |
| 12 | -8 | 7 | $-0.212\ 010\ 620\ 701\ 220 \times 10^4$ | 27 | 0 | 0 | $0.899\ 345\ 518\ 944\ 240$ |
| 13 | -8 | 8 | $0.177\ 683\ 337\ 348\ 191 \times 10^8$ | 28 | 1 | 0 | $-0.211\ 346\ 402\ 240\ 858$ |
| 14 | -6 | 4 | $0.110\ 177\ 443\ 629\ 575 \times 10^2$ | 29 | 1 | 2 | $0.249\ 971\ 752\ 957\ 491 \times 10^2$ |
| 15 | -6 | 6 | $-0.234\ 396\ 091\ 693\ 313 \times 10^6$ | | | | |

Table A1.9. Coefficients and exponents of the backward equation $v_{3i}(p, T)$ for subregion 3i

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|--|-----|-------|-------|--|
| 1 | 0 | 0 | 0.106 905 684 359 136 $\times 10^1$ | 22 | 12 | -12 | 0.164 395 334 345 040 $\times 10^{-23}$ |
| 2 | 0 | 1 | -0.148 620 857 922 333 $\times 10^1$ | 23 | 12 | -6 | -0.339 823 323 754 373 $\times 10^{-5}$ |
| 3 | 0 | 10 | 0.259 862 256 980 408 $\times 10^{15}$ | 24 | 12 | -4 | -0.135 268 639 905 021 $\times 10^{-1}$ |
| 4 | 1 | -4 | -0.446 352 055 678 749 $\times 10^{-11}$ | 25 | 14 | -10 | -0.723 252 514 211 625 $\times 10^{-14}$ |
| 5 | 1 | -2 | -0.566 620 757 170 032 $\times 10^{-6}$ | 26 | 14 | -8 | 0.184 386 437 538 366 $\times 10^{-8}$ |
| 6 | 1 | -1 | -0.235 302 885 736 849 $\times 10^{-2}$ | 27 | 14 | -4 | -0.463 959 533 752 385 $\times 10^{-1}$ |
| 7 | 1 | 0 | -0.269 226 321 968 839 | 28 | 14 | 5 | -0.992 263 100 376 750 $\times 10^{14}$ |
| 8 | 2 | 0 | 0.922 024 992 944 392 $\times 10^1$ | 29 | 18 | -12 | 0.688 169 154 439 335 $\times 10^{-16}$ |
| 9 | 3 | -5 | 0.357 633 505 503 772 $\times 10^{-11}$ | 30 | 18 | -10 | -0.222 620 998 452 197 $\times 10^{-10}$ |
| 10 | 3 | 0 | -0.173 942 565 562 222 $\times 10^2$ | 31 | 18 | -8 | -0.540 843 018 624 083 $\times 10^{-7}$ |
| 11 | 4 | -3 | 0.700 681 785 556 229 $\times 10^{-5}$ | 32 | 18 | -6 | 0.345 570 606 200 257 $\times 10^{-2}$ |
| 12 | 4 | -2 | -0.267 050 351 075 768 $\times 10^{-3}$ | 33 | 18 | 2 | 0.422 275 800 304 086 $\times 10^{11}$ |
| 13 | 4 | -1 | -0.231 779 669 675 624 $\times 10^1$ | 34 | 20 | -12 | -0.126 974 478 770 487 $\times 10^{-14}$ |
| 14 | 5 | -6 | -0.753 533 046 979 752 $\times 10^{-12}$ | 35 | 20 | -10 | 0.927 237 985 153 679 $\times 10^{-9}$ |
| 15 | 5 | -1 | 0.481 337 131 452 891 $\times 10^1$ | 36 | 22 | -12 | 0.612 670 812 016 489 $\times 10^{-13}$ |
| 16 | 5 | 12 | -0.223 286 270 422 356 $\times 10^{22}$ | 37 | 24 | -12 | -0.722 693 924 063 497 $\times 10^{-11}$ |
| 17 | 7 | -4 | -0.118 746 004 987 383 $\times 10^{-4}$ | 38 | 24 | -8 | -0.383 669 502 636 822 $\times 10^{-3}$ |
| 18 | 7 | -3 | 0.646 412 934 136 496 $\times 10^{-2}$ | 39 | 32 | -10 | 0.374 684 572 410 204 $\times 10^{-3}$ |
| 19 | 8 | -6 | -0.410 588 536 330 937 $\times 10^{-9}$ | 40 | 32 | -5 | -0.931 976 897 511 086 $\times 10^5$ |
| 20 | 8 | 10 | 0.422 739 537 057 241 $\times 10^{20}$ | 41 | 36 | -10 | -0.247 690 616 026 922 $\times 10^{-1}$ |
| 21 | 10 | -8 | 0.313 698 180 473 812 $\times 10^{-12}$ | 42 | 36 | -8 | 0.658 110 546 759 474 $\times 10^2$ |

Table A1.10. Coefficients and exponents of the backward equation $v_{3j}(p, T)$ for subregion 3j

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|--|
| 1 | 0 | -1 | -0.111 371 317 395 540 $\times 10^{-3}$ | 16 | 10 | -6 | -0.960 754 116 701 669 $\times 10^{-8}$ |
| 2 | 0 | 0 | 0.100 342 892 423 685 $\times 10^1$ | 17 | 12 | -8 | -0.510 572 269 720 488 $\times 10^{-10}$ |
| 3 | 0 | 1 | 0.530 615 581 928 979 $\times 10^1$ | 18 | 12 | -3 | 0.767 373 781 404 211 $\times 10^{-2}$ |
| 4 | 1 | -2 | 0.179 058 760 078 792 $\times 10^{-5}$ | 19 | 14 | -10 | 0.663 855 469 485 254 $\times 10^{-14}$ |
| 5 | 1 | -1 | -0.728 541 958 464 774 $\times 10^{-3}$ | 20 | 14 | -8 | -0.717 590 735 526 745 $\times 10^{-9}$ |
| 6 | 1 | 1 | -0.187 576 133 371 704 $\times 10^2$ | 21 | 14 | -5 | 0.146 564 542 926 508 $\times 10^{-4}$ |
| 7 | 2 | -1 | 0.199 060 874 071 849 $\times 10^{-2}$ | 22 | 16 | -10 | 0.309 029 474 277 013 $\times 10^{-11}$ |
| 8 | 2 | 1 | 0.243 574 755 377 290 $\times 10^2$ | 23 | 18 | -12 | -0.464 216 300 971 708 $\times 10^{-15}$ |
| 9 | 3 | -2 | -0.177 040 785 499 444 $\times 10^{-3}$ | 24 | 20 | -12 | -0.390 499 637 961 161 $\times 10^{-13}$ |
| 10 | 4 | -2 | -0.259 680 385 227 130 $\times 10^{-2}$ | 25 | 20 | -10 | -0.236 716 126 781 431 $\times 10^{-9}$ |
| 11 | 4 | 2 | -0.198 704 578 406 823 $\times 10^3$ | 26 | 24 | -12 | 0.454 652 854 268 717 $\times 10^{-11}$ |
| 12 | 5 | -3 | 0.738 627 790 224 287 $\times 10^{-4}$ | 27 | 24 | -6 | -0.422 271 787 482 497 $\times 10^{-2}$ |
| 13 | 5 | -2 | -0.236 264 692 844 138 $\times 10^{-2}$ | 28 | 28 | -12 | 0.283 911 742 354 706 $\times 10^{-10}$ |
| 14 | 5 | 0 | -0.161 023 121 314 333 $\times 10^1$ | 29 | 28 | -5 | 0.270 929 002 720 228 $\times 10^1$ |
| 15 | 6 | 3 | 0.622 322 971 786 473 $\times 10^4$ | | | | |

Table A1.11. Coefficients and exponents of the backward equation $v_{3k}(p, T)$ for subregion 3k

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|--|-----|-------|-------|--|
| 1 | -2 | 10 | -0.401 215 699 576 099 $\times 10^9$ | 18 | 1 | 2 | -0.194 646 110 037 079 $\times 10^3$ |
| 2 | -2 | 12 | 0.484 501 478 318 406 $\times 10^{11}$ | 19 | 2 | -8 | 0.808 354 639 772 825 $\times 10^{-15}$ |
| 3 | -1 | -5 | 0.394 721 471 363 678 $\times 10^{-14}$ | 20 | 2 | -6 | -0.180 845 209 145 470 $\times 10^{-10}$ |
| 4 | -1 | 6 | 0.372 629 967 374 147 $\times 10^5$ | 21 | 2 | -3 | -0.696 664 158 132 412 $\times 10^{-5}$ |
| 5 | 0 | -12 | -0.369 794 374 168 666 $\times 10^{-29}$ | 22 | 2 | -2 | -0.181 057 560 300 994 $\times 10^{-2}$ |
| 6 | 0 | -6 | -0.380 436 407 012 452 $\times 10^{-14}$ | 23 | 2 | 0 | 0.255 830 298 579 027 $\times 10^1$ |
| 7 | 0 | -2 | 0.475 361 629 970 233 $\times 10^{-6}$ | 24 | 2 | 4 | 0.328 913 873 658 481 $\times 10^4$ |
| 8 | 0 | -1 | -0.879 148 916 140 706 $\times 10^{-3}$ | 25 | 5 | -12 | -0.173 270 241 249 904 $\times 10^{-18}$ |
| 9 | 0 | 0 | 0.844 317 863 844 331 | 26 | 5 | -6 | -0.661 876 792 558 034 $\times 10^{-6}$ |
| 10 | 0 | 1 | 0.122 433 162 656 600 $\times 10^2$ | 27 | 5 | -3 | -0.395 688 923 421 250 $\times 10^{-2}$ |
| 11 | 0 | 2 | -0.104 529 634 830 279 $\times 10^3$ | 28 | 6 | -12 | 0.604 203 299 819 132 $\times 10^{-17}$ |
| 12 | 0 | 3 | 0.589 702 771 277 429 $\times 10^3$ | 29 | 6 | -10 | -0.400 879 935 920 517 $\times 10^{-13}$ |
| 13 | 0 | 14 | -0.291 026 851 164 444 $\times 10^{14}$ | 30 | 6 | -8 | 0.160 751 107 464 958 $\times 10^{-8}$ |
| 14 | 1 | -3 | 0.170 343 072 841 850 $\times 10^{-5}$ | 31 | 6 | -5 | 0.383 719 409 025 556 $\times 10^{-4}$ |
| 15 | 1 | -2 | -0.277 617 606 975 748 $\times 10^{-3}$ | 32 | 8 | -12 | -0.649 565 446 702 457 $\times 10^{-14}$ |
| 16 | 1 | 0 | -0.344 709 605 486 686 $\times 10^1$ | 33 | 10 | -12 | -0.149 095 328 506 000 $\times 10^{-11}$ |
| 17 | 1 | 1 | 0.221 333 862 447 095 $\times 10^2$ | 34 | 12 | -10 | 0.541 449 377 329 581 $\times 10^{-8}$ |

Table A1.12. Coefficients and exponents of the backward equation $v_{3l}(p, T)$ for subregion 3l

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|---|
| 1 | -12 | 14 | 0.260 702 058 647 537 $\times 10^{10}$ | 23 | -3 | 20 | -0.695 953 622 348 829 $\times 10^{33}$ |
| 2 | -12 | 16 | -0.188 277 213 604 704 $\times 10^{15}$ | 24 | -2 | 2 | 0.110 609 027 472 280 |
| 3 | -12 | 18 | 0.554 923 870 289 667 $\times 10^{19}$ | 25 | -2 | 3 | 0.721 559 163 361 354 $\times 10^2$ |
| 4 | -12 | 20 | -0.758 966 946 387 758 $\times 10^{23}$ | 26 | -2 | 10 | -0.306 367 307 532 219 $\times 10^{15}$ |
| 5 | -12 | 22 | 0.413 865 186 848 908 $\times 10^{27}$ | 27 | -1 | 0 | 0.265 839 618 885 530 $\times 10^{-4}$ |
| 6 | -10 | 14 | -0.815 038 000 738 060 $\times 10^{12}$ | 28 | -1 | 1 | 0.253 392 392 889 754 $\times 10^{-1}$ |
| 7 | -10 | 24 | -0.381 458 260 489 955 $\times 10^{33}$ | 29 | -1 | 3 | -0.214 443 041 836 579 $\times 10^3$ |
| 8 | -8 | 6 | -0.123 239 564 600 519 $\times 10^{-1}$ | 30 | 0 | 0 | 0.937 846 601 489 667 |
| 9 | -8 | 10 | 0.226 095 631 437 174 $\times 10^8$ | 31 | 0 | 1 | 0.223 184 043 101 700 $\times 10^1$ |
| 10 | -8 | 12 | -0.495 017 809 506 720 $\times 10^{12}$ | 32 | 0 | 2 | 0.338 401 222 509 191 $\times 10^2$ |
| 11 | -8 | 14 | 0.529 482 996 422 863 $\times 10^{16}$ | 33 | 0 | 12 | 0.494 237 237 179 718 $\times 10^{21}$ |
| 12 | -8 | 18 | -0.444 359 478 746 295 $\times 10^{23}$ | 34 | 1 | 0 | -0.198 068 404 154 428 |
| 13 | -8 | 24 | 0.521 635 864 527 315 $\times 10^{35}$ | 35 | 1 | 16 | -0.141 415 349 881 140 $\times 10^{31}$ |
| 14 | -8 | 36 | -0.487 095 672 740 742 $\times 10^{55}$ | 36 | 2 | 1 | -0.993 862 421 613 651 $\times 10^2$ |
| 15 | -6 | 8 | -0.714 430 209 937 547 $\times 10^6$ | 37 | 4 | 0 | 0.125 070 534 142 731 $\times 10^3$ |
| 16 | -5 | 4 | 0.127 868 634 615 495 | 38 | 5 | 0 | -0.996 473 529 004 439 $\times 10^3$ |
| 17 | -5 | 5 | -0.100 752 127 917 598 $\times 10^2$ | 39 | 5 | 1 | 0.473 137 909 872 765 $\times 10^5$ |
| 18 | -4 | 7 | 0.777 451 437 960 990 $\times 10^7$ | 40 | 6 | 14 | 0.116 662 121 219 322 $\times 10^{33}$ |
| 19 | -4 | 16 | -0.108 105 480 796 471 $\times 10^{25}$ | 41 | 10 | 4 | -0.315 874 976 271 533 $\times 10^{16}$ |
| 20 | -3 | 1 | -0.357 578 581 169 659 $\times 10^{-5}$ | 42 | 10 | 12 | -0.445 703 369 196 945 $\times 10^{33}$ |
| 21 | -3 | 3 | -0.212 857 169 423 484 $\times 10^1$ | 43 | 14 | 10 | 0.642 794 932 373 694 $\times 10^{33}$ |
| 22 | -3 | 18 | 0.270 706 111 085 238 $\times 10^{30}$ | | | | |

Table A1.13. Coefficients and exponents of the backward equation $v_{3m}(p, T)$ for subregion 3m

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|---|
| 1 | 0 | 0 | 0.811 384 363 481 847 | 21 | 28 | 20 | 0.368 193 926 183 570 $\times 10^{60}$ |
| 2 | 3 | 0 | -0.568 199 310 990 094 $\times 10^4$ | 22 | 2 | 22 | 0.170 215 539 458 936 $\times 10^{18}$ |
| 3 | 8 | 0 | -0.178 657 198 172 556 $\times 10^{11}$ | 23 | 16 | 22 | 0.639 234 909 918 741 $\times 10^{42}$ |
| 4 | 20 | 2 | 0.795 537 657 613 427 $\times 10^{32}$ | 24 | 0 | 24 | -0.821 698 160 721 956 $\times 10^{15}$ |
| 5 | 1 | 5 | -0.814 568 209 346 872 $\times 10^5$ | 25 | 5 | 24 | -0.795 260 241 872 306 $\times 10^{24}$ |
| 6 | 3 | 5 | -0.659 774 567 602 874 $\times 10^8$ | 26 | 0 | 28 | 0.233 415 869 478 510 $\times 10^{18}$ |
| 7 | 4 | 5 | -0.152 861 148 659 302 $\times 10^{11}$ | 27 | 3 | 28 | -0.600 079 934 586 803 $\times 10^{23}$ |
| 8 | 5 | 5 | -0.560 165 667 510 446 $\times 10^{12}$ | 28 | 4 | 28 | 0.594 584 382 273 384 $\times 10^{25}$ |
| 9 | 1 | 6 | 0.458 384 828 593 949 $\times 10^6$ | 29 | 12 | 28 | 0.189 461 279 349 492 $\times 10^{40}$ |
| 10 | 6 | 6 | -0.385 754 000 383 848 $\times 10^{14}$ | 30 | 16 | 28 | -0.810 093 428 842 645 $\times 10^{46}$ |
| 11 | 2 | 7 | 0.453 735 800 004 273 $\times 10^8$ | 31 | 1 | 32 | 0.188 813 911 076 809 $\times 10^{22}$ |
| 12 | 4 | 8 | 0.939 454 935 735 563 $\times 10^{12}$ | 32 | 8 | 32 | 0.111 052 244 098 768 $\times 10^{36}$ |
| 13 | 14 | 8 | 0.266 572 856 432 938 $\times 10^{28}$ | 33 | 14 | 32 | 0.291 133 958 602 503 $\times 10^{46}$ |
| 14 | 2 | 10 | -0.547 578 313 899 097 $\times 10^{10}$ | 34 | 0 | 36 | -0.329 421 923 951 460 $\times 10^{22}$ |
| 15 | 5 | 10 | 0.200 725 701 112 386 $\times 10^{15}$ | 35 | 2 | 36 | -0.137 570 282 536 696 $\times 10^{26}$ |
| 16 | 3 | 12 | 0.185 007 245 563 239 $\times 10^{13}$ | 36 | 3 | 36 | 0.181 508 996 303 902 $\times 10^{28}$ |
| 17 | 0 | 14 | 0.185 135 446 828 337 $\times 10^9$ | 37 | 4 | 36 | -0.346 865 122 768 353 $\times 10^{30}$ |
| 18 | 1 | 14 | -0.170 451 090 076 385 $\times 10^{12}$ | 38 | 8 | 36 | -0.211 961 148 774 260 $\times 10^{38}$ |
| 19 | 1 | 18 | 0.157 890 366 037 614 $\times 10^{15}$ | 39 | 14 | 36 | -0.128 617 899 887 675 $\times 10^{49}$ |
| 20 | 1 | 20 | -0.202 530 509 748 774 $\times 10^{16}$ | 40 | 24 | 36 | 0.479 817 895 699 239 $\times 10^{65}$ |

Table A1.14. Coefficients and exponents of the backward equation $v_{3n}(p, T)$ for subregion 3n

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|--|-----|-------|-------|--|
| 1 | 0 | -12 | 0.280 967 799 943 151 $\times 10^{-38}$ | 21 | 3 | -6 | 0.705 412 100 773 699 $\times 10^{-11}$ |
| 2 | 3 | -12 | 0.614 869 006 573 609 $\times 10^{-30}$ | 22 | 4 | -6 | 0.258 585 887 897 486 $\times 10^{-8}$ |
| 3 | 4 | -12 | 0.582 238 667 048 942 $\times 10^{-27}$ | 23 | 2 | -5 | -0.493 111 362 030 162 $\times 10^{-10}$ |
| 4 | 6 | -12 | 0.390 628 369 238 462 $\times 10^{-22}$ | 24 | 4 | -5 | -0.158 649 699 894 543 $\times 10^{-5}$ |
| 5 | 7 | -12 | 0.821 445 758 255 119 $\times 10^{-20}$ | 25 | 7 | -5 | -0.525 037 427 886 100 |
| 6 | 10 | -12 | 0.402 137 961 842 776 $\times 10^{-14}$ | 26 | 4 | -4 | 0.220 019 901 729 615 $\times 10^{-2}$ |
| 7 | 12 | -12 | 0.651 718 171 878 301 $\times 10^{-12}$ | 27 | 3 | -3 | -0.643 064 132 636 925 $\times 10^{-2}$ |
| 8 | 14 | -12 | -0.211 773 355 803 058 $\times 10^{-7}$ | 28 | 5 | -3 | 0.629 154 149 015 048 $\times 10^2$ |
| 9 | 18 | -12 | 0.264 953 354 380 072 $\times 10^{-2}$ | 29 | 6 | -3 | 0.135 147 318 617 061 $\times 10^3$ |
| 10 | 0 | -10 | -0.135 031 446 451 331 $\times 10^{-31}$ | 30 | 0 | -2 | 0.240 560 808 321 713 $\times 10^{-6}$ |
| 11 | 3 | -10 | -0.607 246 643 970 893 $\times 10^{-23}$ | 31 | 0 | -1 | -0.890 763 306 701 305 $\times 10^{-3}$ |
| 12 | 5 | -10 | -0.402 352 115 234 494 $\times 10^{-18}$ | 32 | 3 | -1 | -0.440 209 599 407 714 $\times 10^4$ |
| 13 | 6 | -10 | -0.744 938 506 925 544 $\times 10^{-16}$ | 33 | 1 | 0 | -0.302 807 107 747 776 $\times 10^3$ |
| 14 | 8 | -10 | 0.189 917 206 526 237 $\times 10^{-12}$ | 34 | 0 | 1 | 0.159 158 748 314 599 $\times 10^4$ |
| 15 | 12 | -10 | 0.364 975 183 508 473 $\times 10^{-5}$ | 35 | 1 | 1 | 0.232 534 272 709 876 $\times 10^6$ |
| 16 | 0 | -8 | 0.177 274 872 361 946 $\times 10^{-25}$ | 36 | 0 | 2 | -0.792 681 207 132 600 $\times 10^6$ |
| 17 | 3 | -8 | -0.334 952 758 812 999 $\times 10^{-18}$ | 37 | 1 | 4 | -0.869 871 364 662 769 $\times 10^{11}$ |
| 18 | 7 | -8 | -0.421 537 726 098 389 $\times 10^{-8}$ | 38 | 0 | 5 | 0.354 542 769 185 671 $\times 10^{12}$ |
| 19 | 12 | -8 | -0.391 048 167 929 649 $\times 10^{-1}$ | 39 | 1 | 6 | 0.400 849 240 129 329 $\times 10^{15}$ |
| 20 | 2 | -6 | 0.541 276 911 564 176 $\times 10^{-13}$ | | | | |

Table A1.15. Coefficients and exponents of the backward equation $v_{3o}(p, T)$ for subregion 3o

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|--|-----|-------|-------|--|
| 1 | 0 | -12 | 0.128 746 023 979 718 $\times 10^{-34}$ | 13 | 6 | -8 | 0.814 897 605 805 513 $\times 10^{-14}$ |
| 2 | 0 | -4 | -0.735 234 770 382 342 $\times 10^{-11}$ | 14 | 7 | -12 | 0.425 596 631 351 839 $\times 10^{-25}$ |
| 3 | 0 | -1 | 0.289 078 692 149 150 $\times 10^{-2}$ | 15 | 8 | -10 | -0.387 449 113 787 755 $\times 10^{-17}$ |
| 4 | 2 | -1 | 0.244 482 731 907 223 | 16 | 8 | -8 | 0.139 814 747 930 240 $\times 10^{-12}$ |
| 5 | 3 | -10 | 0.141 733 492 030 985 $\times 10^{-23}$ | 17 | 8 | -4 | -0.171 849 638 951 521 $\times 10^{-2}$ |
| 6 | 4 | -12 | -0.354 533 853 059 476 $\times 10^{-28}$ | 18 | 10 | -12 | 0.641 890 529 513 296 $\times 10^{-21}$ |
| 7 | 4 | -8 | -0.594 539 202 901 431 $\times 10^{-17}$ | 19 | 10 | -8 | 0.118 960 578 072 018 $\times 10^{-10}$ |
| 8 | 4 | -5 | -0.585 188 401 782 779 $\times 10^{-8}$ | 20 | 14 | -12 | -0.155 282 762 571 611 $\times 10^{-17}$ |
| 9 | 4 | -4 | 0.201 377 325 411 803 $\times 10^{-5}$ | 21 | 14 | -8 | 0.233 907 907 347 507 $\times 10^{-7}$ |
| 10 | 4 | -1 | 0.138 647 388 209 306 $\times 10^1$ | 22 | 20 | -12 | -0.174 093 247 766 213 $\times 10^{-12}$ |
| 11 | 5 | -4 | -0.173 959 365 084 772 $\times 10^{-4}$ | 23 | 20 | -10 | 0.377 682 649 089 149 $\times 10^{-8}$ |
| 12 | 5 | -3 | 0.137 680 878 349 369 $\times 10^{-2}$ | 24 | 24 | -12 | -0.516 720 236 575 302 $\times 10^{-10}$ |

Table A1.16. Coefficients and exponents of the backward equation $v_{3p}(p, T)$ for subregion 3p

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|--|
| 1 | 0 | -1 | -0.982 825 342 010 366 $\times 10^{-4}$ | 15 | 12 | -12 | 0.343 480 022 104 968 $\times 10^{-25}$ |
| 2 | 0 | 0 | 0.105 145 700 850 612 $\times 10^1$ | 16 | 12 | -6 | 0.816 256 095 947 021 $\times 10^{-5}$ |
| 3 | 0 | 1 | 0.116 033 094 095 084 $\times 10^3$ | 17 | 12 | -5 | 0.294 985 697 916 798 $\times 10^{-2}$ |
| 4 | 0 | 2 | 0.324 664 750 281 543 $\times 10^4$ | 18 | 14 | -10 | 0.711 730 466 276 584 $\times 10^{-16}$ |
| 5 | 1 | 1 | -0.123 592 348 610 137 $\times 10^4$ | 19 | 14 | -8 | 0.400 954 763 806 941 $\times 10^{-9}$ |
| 6 | 2 | -1 | -0.561 403 450 013 495 $\times 10^{-1}$ | 20 | 14 | -3 | 0.107 766 027 032 853 $\times 10^2$ |
| 7 | 3 | -3 | 0.856 677 401 640 869 $\times 10^{-7}$ | 21 | 16 | -8 | -0.409 449 599 138 182 $\times 10^{-6}$ |
| 8 | 3 | 0 | 0.236 313 425 393 924 $\times 10^3$ | 22 | 18 | -8 | -0.729 121 307 758 902 $\times 10^{-5}$ |
| 9 | 4 | -2 | 0.972 503 292 350 109 $\times 10^{-2}$ | 23 | 20 | -10 | 0.677 107 970 938 909 $\times 10^{-8}$ |
| 10 | 6 | -2 | -0.103 001 994 531 927 $\times 10^1$ | 24 | 22 | -10 | 0.602 745 973 022 975 $\times 10^{-7}$ |
| 11 | 7 | -5 | -0.149 653 706 199 162 $\times 10^{-8}$ | 25 | 24 | -12 | -0.382 323 011 855 257 $\times 10^{-10}$ |
| 12 | 7 | -4 | -0.215 743 778 861 592 $\times 10^{-4}$ | 26 | 24 | -8 | 0.179 946 628 317 437 $\times 10^{-2}$ |
| 13 | 8 | -2 | -0.834 452 198 291 445 $\times 10^1$ | 27 | 36 | -12 | -0.345 042 834 640 005 $\times 10^{-3}$ |
| 14 | 10 | -3 | 0.586 602 660 564 988 | | | | |

Table A1.17. Coefficients and exponents of the backward equation $v_{3q}(p, T)$ for subregion 3q

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|---|
| 1 | -12 | 10 | -0.820 433 843 259 950 $\times 10^5$ | 13 | -3 | 3 | 0.232 808 472 983 776 $\times 10^3$ |
| 2 | -12 | 12 | 0.473 271 518 461 586 $\times 10^{11}$ | 14 | -2 | 0 | -0.142 808 220 416 837 $\times 10^{-4}$ |
| 3 | -10 | 6 | -0.805 950 021 005 413 $\times 10^{-1}$ | 15 | -2 | 1 | -0.643 596 060 678 456 $\times 10^{-2}$ |
| 4 | -10 | 7 | 0.328 600 025 435 980 $\times 10^2$ | 16 | -2 | 2 | -0.428 577 227 475 614 $\times 10^1$ |
| 5 | -10 | 8 | -0.356 617 029 982 490 $\times 10^4$ | 17 | -2 | 4 | 0.225 689 939 161 918 $\times 10^4$ |
| 6 | -10 | 10 | -0.172 985 781 433 335 $\times 10^{10}$ | 18 | -1 | 0 | 0.100 355 651 721 510 $\times 10^{-2}$ |
| 7 | -8 | 8 | 0.351 769 232 729 192 $\times 10^8$ | 19 | -1 | 1 | 0.333 491 455 143 516 |
| 8 | -6 | 6 | -0.775 489 259 985 144 $\times 10^6$ | 20 | -1 | 2 | 0.109 697 576 888 873 $\times 10^1$ |
| 9 | -5 | 2 | 0.710 346 691 966 018 $\times 10^{-4}$ | 21 | 0 | 0 | 0.961 917 379 376 452 |
| 10 | -5 | 5 | 0.993 499 883 820 274 $\times 10^5$ | 22 | 1 | 0 | -0.838 165 632 204 598 $\times 10^{-1}$ |
| 11 | -4 | 3 | -0.642 094 171 904 570 | 23 | 1 | 1 | 0.247 795 908 411 492 $\times 10^1$ |
| 12 | -4 | 4 | -0.612 842 816 820 083 $\times 10^4$ | 24 | 1 | 3 | -0.319 114 969 006 533 $\times 10^4$ |

Table A1.18. Coefficients and exponents of the backward equation $v_{3r}(p, T)$ for subregion 3r

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|--|-----|-------|-------|---|
| 1 | -8 | 6 | 0.144 165 955 660 863 $\times 10^{-2}$ | 15 | 8 | -10 | 0.399 988 795 693 162 $\times 10^{-12}$ |
| 2 | -8 | 14 | -0.701 438 599 628 258 $\times 10^{13}$ | 16 | 8 | -8 | -0.536 479 560 201 811 $\times 10^{-6}$ |
| 3 | -3 | -3 | -0.830 946 716 459 219 $\times 10^{-16}$ | 17 | 8 | -5 | 0.159 536 722 411 202 $\times 10^{-1}$ |
| 4 | -3 | 3 | 0.261 975 135 368 109 | 18 | 10 | -12 | 0.270 303 248 860 217 $\times 10^{-14}$ |
| 5 | -3 | 4 | 0.393 097 214 706 245 $\times 10^3$ | 19 | 10 | -10 | 0.244 247 453 858 506 $\times 10^{-7}$ |
| 6 | -3 | 5 | -0.104 334 030 654 021 $\times 10^5$ | 20 | 10 | -8 | -0.983 430 636 716 454 $\times 10^{-5}$ |
| 7 | -3 | 8 | 0.490 112 654 154 211 $\times 10^9$ | 21 | 10 | -6 | 0.663 513 144 224 454 $\times 10^{-1}$ |
| 8 | 0 | -1 | -0.147 104 222 772 069 $\times 10^{-3}$ | 22 | 10 | -5 | -0.993 456 957 845 006 $\times 10^1$ |
| 9 | 0 | 0 | 0.103 602 748 043 408 $\times 10^1$ | 23 | 10 | -4 | 0.546 491 323 528 491 $\times 10^3$ |
| 10 | 0 | 1 | 0.305 308 890 065 089 $\times 10^1$ | 24 | 10 | -3 | -0.143 365 406 393 758 $\times 10^5$ |
| 11 | 0 | 5 | -0.399 745 276 971 264 $\times 10^7$ | 25 | 10 | -2 | 0.150 764 974 125 511 $\times 10^6$ |
| 12 | 3 | -6 | 0.569 233 719 593 750 $\times 10^{-11}$ | 26 | 12 | -12 | -0.337 209 709 340 105 $\times 10^{-9}$ |
| 13 | 3 | -2 | -0.464 923 504 407 778 $\times 10^{-1}$ | 27 | 14 | -12 | 0.377 501 980 025 469 $\times 10^{-8}$ |
| 14 | 8 | -12 | -0.535 400 396 512 906 $\times 10^{-17}$ | | | | |

Table A1.19. Coefficients and exponents of the backward equation $v_{3s}(p, T)$ for subregion 3s

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|---|
| 1 | -12 | 20 | -0.532 466 612 140 254 $\times 10^{23}$ | 16 | 0 | 0 | 0.965 961 650 599 775 |
| 2 | -12 | 24 | 0.100 415 480 000 824 $\times 10^{32}$ | 17 | 0 | 1 | 0.294 885 696 802 488 $\times 10^1$ |
| 3 | -10 | 22 | -0.191 540 001 821 367 $\times 10^{30}$ | 18 | 0 | 4 | -0.653 915 627 346 115 $\times 10^5$ |
| 4 | -8 | 14 | 0.105 618 377 808 847 $\times 10^{17}$ | 19 | 0 | 28 | 0.604 012 200 163 444 $\times 10^{50}$ |
| 5 | -6 | 36 | 0.202 281 884 477 061 $\times 10^{59}$ | 20 | 1 | 0 | -0.198 339 358 557 937 |
| 6 | -5 | 8 | 0.884 585 472 596 134 $\times 10^8$ | 21 | 1 | 32 | -0.175 984 090 163 501 $\times 10^{58}$ |
| 7 | -5 | 16 | 0.166 540 181 638 363 $\times 10^{23}$ | 22 | 3 | 0 | 0.356 314 881 403 987 $\times 10^1$ |
| 8 | -4 | 6 | -0.313 563 197 669 111 $\times 10^6$ | 23 | 3 | 1 | -0.575 991 255 144 384 $\times 10^3$ |
| 9 | -4 | 32 | -0.185 662 327 545 324 $\times 10^{54}$ | 24 | 3 | 2 | 0.456 213 415 338 071 $\times 10^5$ |
| 10 | -3 | 3 | -0.624 942 093 918 942 $\times 10^{-1}$ | 25 | 4 | 3 | -0.109 174 044 987 829 $\times 10^8$ |
| 11 | -3 | 8 | -0.504 160 724 132 590 $\times 10^{10}$ | 26 | 4 | 18 | 0.437 796 099 975 134 $\times 10^{34}$ |
| 12 | -2 | 4 | 0.187 514 491 833 092 $\times 10^5$ | 27 | 4 | 24 | -0.616 552 611 135 792 $\times 10^{46}$ |
| 13 | -1 | 1 | 0.121 399 979 993 217 $\times 10^{-2}$ | 28 | 5 | 4 | 0.193 568 768 917 797 $\times 10^{10}$ |
| 14 | -1 | 2 | 0.188 317 043 049 455 $\times 10^1$ | 29 | 14 | 24 | 0.950 898 170 425 042 $\times 10^{54}$ |
| 15 | -1 | 3 | -0.167 073 503 962 060 $\times 10^4$ | | | | |

Table A1.20. Coefficients and exponents of the backward equation $v_{3t}(p, T)$ for subregion 3t

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|---|
| 1 | 0 | 0 | $0.155\ 287\ 249\ 586\ 268 \times 10^1$ | 18 | 7 | 36 | $-0.341\ 552\ 040\ 860\ 644 \times 10^{51}$ |
| 2 | 0 | 1 | $0.664\ 235\ 115\ 009\ 031 \times 10^1$ | 19 | 10 | 10 | $-0.527\ 251\ 339\ 709\ 047 \times 10^{21}$ |
| 3 | 0 | 4 | $-0.289\ 366\ 236\ 727\ 210 \times 10^4$ | 20 | 10 | 12 | $0.245\ 375\ 640\ 937\ 055 \times 10^{24}$ |
| 4 | 0 | 12 | $-0.385\ 923\ 202\ 309\ 848 \times 10^{13}$ | 21 | 10 | 14 | $-0.168\ 776\ 617\ 209\ 269 \times 10^{27}$ |
| 5 | 1 | 0 | $-0.291\ 002\ 915\ 783\ 761 \times 10^1$ | 22 | 10 | 16 | $0.358\ 958\ 955\ 867\ 578 \times 10^{29}$ |
| 6 | 1 | 10 | $-0.829\ 088\ 246\ 858\ 083 \times 10^{12}$ | 23 | 10 | 22 | $-0.656\ 475\ 280\ 339\ 411 \times 10^{36}$ |
| 7 | 2 | 0 | $0.176\ 814\ 899\ 675\ 218 \times 10^1$ | 24 | 18 | 18 | $0.355\ 286\ 045\ 512\ 301 \times 10^{39}$ |
| 8 | 2 | 6 | $-0.534\ 686\ 695\ 713\ 469 \times 10^9$ | 25 | 20 | 32 | $0.569\ 021\ 454\ 413\ 270 \times 10^{58}$ |
| 9 | 2 | 14 | $0.160\ 464\ 608\ 687\ 834 \times 10^{18}$ | 26 | 22 | 22 | $-0.700\ 584\ 546\ 433\ 113 \times 10^{48}$ |
| 10 | 3 | 3 | $0.196\ 435\ 366\ 560\ 186 \times 10^6$ | 27 | 22 | 36 | $-0.705\ 772\ 623\ 326\ 374 \times 10^{65}$ |
| 11 | 3 | 8 | $0.156\ 637\ 427\ 541\ 729 \times 10^{13}$ | 28 | 24 | 24 | $0.166\ 861\ 176\ 200\ 148 \times 10^{53}$ |
| 12 | 4 | 0 | $-0.178\ 154\ 560\ 260\ 006 \times 10^1$ | 29 | 28 | 28 | $-0.300\ 475\ 129\ 680\ 486 \times 10^{61}$ |
| 13 | 4 | 10 | $-0.229\ 746\ 237\ 623\ 692 \times 10^{16}$ | 30 | 32 | 22 | $-0.668\ 481\ 295\ 196\ 808 \times 10^{51}$ |
| 14 | 7 | 3 | $0.385\ 659\ 001\ 648\ 006 \times 10^8$ | 31 | 32 | 32 | $0.428\ 432\ 338\ 620\ 678 \times 10^{69}$ |
| 15 | 7 | 4 | $0.110\ 554\ 446\ 790\ 543 \times 10^{10}$ | 32 | 32 | 36 | $-0.444\ 227\ 367\ 758\ 304 \times 10^{72}$ |
| 16 | 7 | 7 | $-0.677\ 073\ 830\ 687\ 349 \times 10^{14}$ | 33 | 36 | 36 | $-0.281\ 396\ 013\ 562\ 745 \times 10^{77}$ |
| 17 | 7 | 20 | $-0.327\ 910\ 592\ 086\ 523 \times 10^{31}$ | | | | |

A2 Coefficients for Auxiliary Equations**Table A2.1.** Coefficients and exponents of the auxiliary equation $v_{3u}(p, T)$ for subregion 3u

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|--|
| 1 | -12 | 14 | $0.122\ 088\ 349\ 258\ 355 \times 10^{18}$ | 20 | 1 | -2 | $0.105\ 581\ 745\ 346\ 187 \times 10^{-2}$ |
| 2 | -10 | 10 | $0.104\ 216\ 468\ 608\ 488 \times 10^{10}$ | 21 | 2 | 5 | $-0.651\ 903\ 203\ 602\ 581 \times 10^{15}$ |
| 3 | -10 | 12 | $-0.882\ 666\ 931\ 564\ 652 \times 10^{16}$ | 22 | 2 | 10 | $-0.160\ 116\ 813\ 274\ 676 \times 10^{25}$ |
| 4 | -10 | 14 | $0.259\ 929\ 510\ 849\ 499 \times 10^{20}$ | 23 | 3 | -5 | $-0.510\ 254\ 294\ 237\ 837 \times 10^{-8}$ |
| 5 | -8 | 10 | $0.222\ 612\ 779\ 142\ 211 \times 10^{15}$ | 24 | 5 | -4 | $-0.152\ 355\ 388\ 953\ 402$ |
| 6 | -8 | 12 | $-0.878\ 473\ 585\ 050\ 085 \times 10^{18}$ | 25 | 5 | 2 | $0.677\ 143\ 292\ 290\ 144 \times 10^{12}$ |
| 7 | -8 | 14 | $-0.314\ 432\ 577\ 551\ 552 \times 10^{22}$ | 26 | 5 | 3 | $0.276\ 378\ 438\ 378\ 930 \times 10^{15}$ |
| 8 | -6 | 8 | $-0.216\ 934\ 916\ 996\ 285 \times 10^{13}$ | 27 | 6 | -5 | $0.116\ 862\ 983\ 141\ 686 \times 10^{-1}$ |
| 9 | -6 | 12 | $0.159\ 079\ 648\ 196\ 849 \times 10^{21}$ | 28 | 6 | 2 | $-0.301\ 426\ 947\ 980\ 171 \times 10^{14}$ |
| 10 | -5 | 4 | $-0.339\ 567\ 617\ 303\ 423 \times 10^3$ | 29 | 8 | -8 | $0.169\ 719\ 813\ 884\ 840 \times 10^{-7}$ |
| 11 | -5 | 8 | $0.884\ 387\ 651\ 337\ 836 \times 10^{13}$ | 30 | 8 | 8 | $0.104\ 674\ 840\ 020\ 929 \times 10^{27}$ |
| 12 | -5 | 12 | $-0.843\ 405\ 926\ 846\ 418 \times 10^{21}$ | 31 | 10 | -4 | $-0.108\ 016\ 904\ 560\ 140 \times 10^5$ |
| 13 | -3 | 2 | $0.114\ 178\ 193\ 518\ 022 \times 10^2$ | 32 | 12 | -12 | $-0.990\ 623\ 601\ 934\ 295 \times 10^{-12}$ |
| 14 | -1 | -1 | $-0.122\ 708\ 229\ 235\ 641 \times 10^{-3}$ | 33 | 12 | -4 | $0.536\ 116\ 483\ 602\ 738 \times 10^7$ |
| 15 | -1 | 1 | $-0.106\ 201\ 671\ 767\ 107 \times 10^3$ | 34 | 12 | 4 | $0.226\ 145\ 963\ 747\ 881 \times 10^{22}$ |
| 16 | -1 | 12 | $0.903\ 443\ 213\ 959\ 313 \times 10^{25}$ | 35 | 14 | -12 | $-0.488\ 731\ 565\ 776\ 210 \times 10^{-9}$ |
| 17 | -1 | 14 | $-0.693\ 996\ 270\ 370\ 852 \times 10^{28}$ | 36 | 14 | -10 | $0.151\ 001\ 548\ 880\ 670 \times 10^{-4}$ |
| 18 | 0 | -3 | $0.648\ 916\ 718\ 965\ 575 \times 10^{-8}$ | 37 | 14 | -6 | $-0.227\ 700\ 464\ 643\ 920 \times 10^5$ |
| 19 | 0 | 1 | $0.718\ 957\ 567\ 127\ 851 \times 10^4$ | 38 | 14 | 6 | $-0.781\ 754\ 507\ 698\ 846 \times 10^{28}$ |

Table A2.2. Coefficients and exponents of the auxiliary equation $v_{3v}(p, T)$ for subregion 3v

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|--|-----|-------|-------|--|
| 1 | -10 | -8 | -0.415 652 812 061 591 $\times 10^{-54}$ | 21 | -3 | 12 | 0.742 705 723 302 738 $\times 10^{27}$ |
| 2 | -8 | -12 | 0.177 441 742 924 043 $\times 10^{-60}$ | 22 | -2 | 2 | -0.517 429 682 450 605 $\times 10^2$ |
| 3 | -6 | -12 | -0.357 078 668 203 377 $\times 10^{-54}$ | 23 | -2 | 4 | 0.820 612 048 645 469 $\times 10^7$ |
| 4 | -6 | -3 | 0.359 252 213 604 114 $\times 10^{-25}$ | 24 | -1 | -2 | -0.188 214 882 341 448 $\times 10^{-8}$ |
| 5 | -6 | 5 | -0.259 123 736 380 269 $\times 10^2$ | 25 | -1 | 0 | 0.184 587 261 114 837 $\times 10^{-1}$ |
| 6 | -6 | 6 | 0.594 619 766 193 460 $\times 10^5$ | 26 | 0 | -2 | -0.135 830 407 782 663 $\times 10^{-5}$ |
| 7 | -6 | 8 | -0.624 184 007 103 158 $\times 10^{11}$ | 27 | 0 | 6 | -0.723 681 885 626 348 $\times 10^{17}$ |
| 8 | -6 | 10 | 0.313 080 299 915 944 $\times 10^{17}$ | 28 | 0 | 10 | -0.223 449 194 054 124 $\times 10^{27}$ |
| 9 | -5 | 1 | 0.105 006 446 192 036 $\times 10^{-8}$ | 29 | 1 | -12 | -0.111 526 741 826 431 $\times 10^{-34}$ |
| 10 | -5 | 2 | -0.192 824 336 984 852 $\times 10^{-5}$ | 30 | 1 | -10 | 0.276 032 601 145 151 $\times 10^{-28}$ |
| 11 | -5 | 6 | 0.654 144 373 749 937 $\times 10^6$ | 31 | 3 | 3 | 0.134 856 491 567 853 $\times 10^{15}$ |
| 12 | -5 | 8 | 0.513 117 462 865 044 $\times 10^{13}$ | 32 | 4 | -6 | 0.652 440 293 345 860 $\times 10^{-9}$ |
| 13 | -5 | 10 | -0.697 595 750 347 391 $\times 10^{19}$ | 33 | 4 | 3 | 0.510 655 119 774 360 $\times 10^{17}$ |
| 14 | -5 | 14 | -0.103 977 184 454 767 $\times 10^{29}$ | 34 | 4 | 10 | -0.468 138 358 908 732 $\times 10^{32}$ |
| 15 | -4 | -12 | 0.119 563 135 540 666 $\times 10^{-47}$ | 35 | 5 | 2 | -0.760 667 491 183 279 $\times 10^{16}$ |
| 16 | -4 | -10 | -0.436 677 034 051 655 $\times 10^{-41}$ | 36 | 8 | -12 | -0.417 247 986 986 821 $\times 10^{-18}$ |
| 17 | -4 | -6 | 0.926 990 036 530 639 $\times 10^{-29}$ | 37 | 10 | -2 | 0.312 545 677 756 104 $\times 10^{14}$ |
| 18 | -4 | 10 | 0.587 793 105 620 748 $\times 10^{21}$ | 38 | 12 | -3 | -0.100 375 333 864 186 $\times 10^{15}$ |
| 19 | -3 | -3 | 0.280 375 725 094 731 $\times 10^{-17}$ | 39 | 14 | 1 | 0.247 761 392 329 058 $\times 10^{27}$ |
| 20 | -3 | 10 | -0.192 359 972 440 634 $\times 10^{23}$ | | | | |

Table A2.3. Coefficients and exponents of the auxiliary equation $v_{3w}(p, T)$ for subregion 3w

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|--|-----|-------|-------|--|
| 1 | -12 | 8 | -0.586 219 133 817 016 $\times 10^{-7}$ | 19 | -1 | -8 | 0.237 416 732 616 644 $\times 10^{-26}$ |
| 2 | -12 | 14 | -0.894 460 355 005 526 $\times 10^{11}$ | 20 | -1 | -4 | 0.271 700 235 739 893 $\times 10^{-14}$ |
| 3 | -10 | -1 | 0.531 168 037 519 774 $\times 10^{-30}$ | 21 | -1 | 1 | -0.907 886 213 483 600 $\times 10^2$ |
| 4 | -10 | 8 | 0.109 892 402 329 239 | 22 | 0 | -12 | -0.171 242 509 570 207 $\times 10^{-36}$ |
| 5 | -8 | 6 | -0.575 368 389 425 212 $\times 10^{-1}$ | 23 | 0 | 1 | 0.156 792 067 854 621 $\times 10^3$ |
| 6 | -8 | 8 | 0.228 276 853 990 249 $\times 10^5$ | 24 | 1 | -1 | 0.923 261 357 901 470 |
| 7 | -8 | 14 | -0.158 548 609 655 002 $\times 10^{19}$ | 25 | 2 | -1 | -0.597 865 988 422 577 $\times 10^1$ |
| 8 | -6 | -4 | 0.329 865 748 576 503 $\times 10^{-27}$ | 26 | 2 | 2 | 0.321 988 767 636 389 $\times 10^7$ |
| 9 | -6 | -3 | -0.634 987 981 190 669 $\times 10^{-24}$ | 27 | 3 | -12 | -0.399 441 390 042 203 $\times 10^{-29}$ |
| 10 | -6 | 2 | 0.615 762 068 640 611 $\times 10^{-8}$ | 28 | 3 | -5 | 0.493 429 086 046 981 $\times 10^{-7}$ |
| 11 | -6 | 8 | -0.961 109 240 985 747 $\times 10^8$ | 29 | 5 | -10 | 0.812 036 983 370 565 $\times 10^{-19}$ |
| 12 | -5 | -10 | -0.406 274 286 652 625 $\times 10^{-44}$ | 30 | 5 | -8 | -0.207 610 284 654 137 $\times 10^{-11}$ |
| 13 | -4 | -1 | -0.471 103 725 498 077 $\times 10^{-12}$ | 31 | 5 | -6 | -0.340 821 291 419 719 $\times 10^{-6}$ |
| 14 | -4 | 3 | 0.725 937 724 828 145 | 32 | 8 | -12 | 0.542 000 573 372 233 $\times 10^{-17}$ |
| 15 | -3 | -10 | 0.187 768 525 763 682 $\times 10^{-38}$ | 33 | 8 | -10 | -0.856 711 586 510 214 $\times 10^{-12}$ |
| 16 | -3 | 3 | -0.103 308 436 323 771 $\times 10^4$ | 34 | 10 | -12 | 0.266 170 454 405 981 $\times 10^{-13}$ |
| 17 | -2 | 1 | -0.662 552 816 342 168 $\times 10^{-1}$ | 35 | 10 | -8 | 0.858 133 791 857 099 $\times 10^{-5}$ |
| 18 | -2 | 2 | 0.579 514 041 765 710 $\times 10^3$ | | | | |

Table A2.4. Coefficients and exponents of the auxiliary equation $v_{3x}(p, T)$ for subregion 3x

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|--|-----|-------|-------|---|
| 1 | -8 | 14 | 0.377 373 741 298 151 $\times 10^{19}$ | 19 | 4 | 3 | 0.397 949 001 553 184 $\times 10^{14}$ |
| 2 | -6 | 10 | -0.507 100 883 722 913 $\times 10^{13}$ | 20 | 5 | -6 | 0.100 824 008 584 757 $\times 10^{-6}$ |
| 3 | -5 | 10 | -0.103 363 225 598 860 $\times 10^{16}$ | 21 | 5 | -2 | 0.162 234 569 738 433 $\times 10^5$ |
| 4 | -4 | 1 | 0.184 790 814 320 773 $\times 10^{-5}$ | 22 | 5 | 1 | -0.432 355 225 319 745 $\times 10^{11}$ |
| 5 | -4 | 2 | -0.924 729 378 390 945 $\times 10^{-3}$ | 23 | 6 | 1 | -0.592 874 245 598 610 $\times 10^{12}$ |
| 6 | -4 | 14 | -0.425 999 562 292 738 $\times 10^{24}$ | 24 | 8 | -6 | 0.133 061 647 281 106 $\times 10^1$ |
| 7 | -3 | -2 | -0.462 307 771 873 973 $\times 10^{-12}$ | 25 | 8 | -3 | 0.157 338 197 797 544 $\times 10^7$ |
| 8 | -3 | 12 | 0.107 319 065 855 767 $\times 10^{22}$ | 26 | 8 | 1 | 0.258 189 614 270 853 $\times 10^{14}$ |
| 9 | -1 | 5 | 0.648 662 492 280 682 $\times 10^{11}$ | 27 | 8 | 8 | 0.262 413 209 706 358 $\times 10^{25}$ |
| 10 | 0 | 0 | 0.244 200 600 688 281 $\times 10^1$ | 28 | 10 | -8 | -0.920 011 937 431 142 $\times 10^{-1}$ |
| 11 | 0 | 4 | -0.851 535 733 484 258 $\times 10^{10}$ | 29 | 12 | -10 | 0.220 213 765 905 426 $\times 10^{-2}$ |
| 12 | 0 | 10 | 0.169 894 481 433 592 $\times 10^{22}$ | 30 | 12 | -8 | -0.110 433 759 109 547 $\times 10^2$ |
| 13 | 1 | -10 | 0.215 780 222 509 020 $\times 10^{-26}$ | 31 | 12 | -5 | 0.847 004 870 612 087 $\times 10^7$ |
| 14 | 1 | -1 | -0.320 850 551 367 334 | 32 | 12 | -4 | -0.592 910 695 762 536 $\times 10^9$ |
| 15 | 2 | 6 | -0.382 642 448 458 610 $\times 10^{17}$ | 33 | 14 | -12 | -0.183 027 173 269 660 $\times 10^{-4}$ |
| 16 | 3 | -12 | -0.275 386 077 674 421 $\times 10^{-28}$ | 34 | 14 | -10 | 0.181 339 603 516 302 |
| 17 | 3 | 0 | -0.563 199 253 391 666 $\times 10^6$ | 35 | 14 | -8 | -0.119 228 759 669 889 $\times 10^4$ |
| 18 | 3 | 8 | -0.326 068 646 279 314 $\times 10^{21}$ | 36 | 14 | -6 | 0.430 867 658 061 468 $\times 10^7$ |

Table A2.5. Coefficients and exponents of the auxiliary equation $v_{3y}(p, T)$ for subregion 3y

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|---|-----|-------|-------|---|
| 1 | 0 | -3 | -0.525 597 995 024 633 $\times 10^{-9}$ | 11 | 3 | 4 | 0.705 106 224 399 834 $\times 10^{21}$ |
| 2 | 0 | 1 | 0.583 441 305 228 407 $\times 10^4$ | 12 | 3 | 8 | -0.266 713 136 106 469 $\times 10^{31}$ |
| 3 | 0 | 5 | -0.134 778 968 457 925 $\times 10^{17}$ | 13 | 4 | -6 | -0.145 370 512 554 562 $\times 10^{-7}$ |
| 4 | 0 | 8 | 0.118 973 500 934 212 $\times 10^{26}$ | 14 | 4 | 6 | 0.149 333 917 053 130 $\times 10^{28}$ |
| 5 | 1 | 8 | -0.159 096 490 904 708 $\times 10^{27}$ | 15 | 5 | -2 | -0.149 795 620 287 641 $\times 10^8$ |
| 6 | 2 | -4 | -0.315 839 902 302 021 $\times 10^{-6}$ | 16 | 5 | 1 | -0.381 881 906 271 100 $\times 10^{16}$ |
| 7 | 2 | -1 | 0.496 212 197 158 239 $\times 10^3$ | 17 | 8 | -8 | 0.724 660 165 585 797 $\times 10^{-4}$ |
| 8 | 2 | 4 | 0.327 777 227 273 171 $\times 10^{19}$ | 18 | 8 | -2 | -0.937 808 169 550 193 $\times 10^{14}$ |
| 9 | 2 | 5 | -0.527 114 657 850 696 $\times 10^{22}$ | 19 | 10 | -5 | 0.514 411 468 376 383 $\times 10^{10}$ |
| 10 | 3 | -8 | 0.210 017 506 281 863 $\times 10^{-16}$ | 20 | 12 | -8 | -0.828 198 594 040 141 $\times 10^5$ |

Table A2.6. Coefficients and exponents of the auxiliary equation $v_{3z}(p, T)$ for subregion 3z

| i | I_i | J_i | n_i | i | I_i | J_i | n_i |
|-----|-------|-------|--|-----|-------|-------|--|
| 1 | -8 | 3 | 0.244 007 892 290 650 $\times 10^{-10}$ | 13 | 0 | 3 | 0.328 380 587 890 663 $\times 10^{12}$ |
| 2 | -6 | 6 | -0.463 057 430 331 242 $\times 10^7$ | 14 | 1 | 1 | -0.625 004 791 171 543 $\times 10^8$ |
| 3 | -5 | 6 | 0.728 803 274 777 712 $\times 10^{10}$ | 15 | 2 | 6 | 0.803 197 957 462 023 $\times 10^{21}$ |
| 4 | -5 | 8 | 0.327 776 302 858 856 $\times 10^{16}$ | 16 | 3 | -6 | -0.204 397 011 338 353 $\times 10^{-10}$ |
| 5 | -4 | 5 | -0.110 598 170 118 409 $\times 10^{10}$ | 17 | 3 | -2 | -0.378 391 047 055 938 $\times 10^4$ |
| 6 | -4 | 6 | -0.323 899 915 729 957 $\times 10^{13}$ | 18 | 6 | -6 | 0.972 876 545 938 620 $\times 10^{-2}$ |
| 7 | -4 | 8 | 0.923 814 007 023 245 $\times 10^{16}$ | 19 | 6 | -5 | 0.154 355 721 681 459 $\times 10^2$ |
| 8 | -3 | -2 | 0.842 250 080 413 712 $\times 10^{-12}$ | 20 | 6 | -4 | -0.373 962 862 928 643 $\times 10^4$ |
| 9 | -3 | 5 | 0.663 221 436 245 506 $\times 10^{12}$ | 21 | 6 | -1 | -0.682 859 011 374 572 $\times 10^{11}$ |
| 10 | -3 | 6 | -0.167 170 186 672 139 $\times 10^{15}$ | 22 | 8 | -8 | -0.248 488 015 614 543 $\times 10^{-3}$ |
| 11 | -2 | 2 | 0.253 749 358 701 391 $\times 10^4$ | 23 | 8 | -4 | 0.394 536 049 497 068 $\times 10^7$ |
| 12 | -1 | -6 | -0.819 731 559 610 523 $\times 10^{-20}$ | | | | |